Does Access Location Influence Outcome of Percutaneous Nephrolithotomy in Staghorn Stones?

Serkan Yarimoglu¹, Murat Sahan¹, Salih Polat², Gurkan Cesur¹, Omer Koras³ and Tansu Degirmenci¹

¹Department of Urology, HSU Izmir Bozyaka Training and Research Hospital, Izmir, Turkey

²Department of Urology, Faculty of Medicine, Amasya University, Amasya, Turkey

³Department of Urology, Faculty of Medicine, Hatay University, Hatay, Turkey

ABSTRACT

Objective: To assess the effect of lower and middle calyceal accesses on the outcomes of percutaneous nephrolithotomy (PCNL) in staghorn stones.

Study Design: Observational study.

Place and Duration of Study: Department of Urology, University of Health Sciences, Izmir Bozyaka Training and Research Hospital, Turkey, from April 2012 to January 2019.

Methodology: Patients who underwent PCNL for staghorn stones were retrospectively analysed. The patients were grouped as Group-1 (middle calyceal access) and Group-2 (lower calyceal access). Demographic and stone characteristics (size, density), perioperative and postoperative data, and stone-free rates were compared between these two groups. Postoperative detection of > 4 mm stones was defined as residual stones.

Results: There were 249 patients in the study; 108 in middle calyceal access group and 141 in lower calyceal access group. The mean stone burden was 765 (524-1322) and 777 (490-1445) mm² in group-1 and group-2, respectively (p=0.876). The number of stone-free patients was 50 (46.3%) in middle calyceal access group and 93 (66.0%) in lower calyceal access group. The rate of stone-free status was significantly higher in lower calyceal access group (p=0.002). The overall complication rate was similar between the groups (p=0.132). The binary analysis showed that stone burden, and calyx of entry were predictive factors for success.

Conclusion: Although the choice of the calyx to be entered does not affect the complication rate in staghorn stones, the stone-free rate is significantly higher in lower calyceal access.

Key Words: Percutaneous nephrolithotomy, Staghorn stones, Clavien scoring system, Calyx, Access.

How to cite this article: Yarimoglu S, Sahan M, Polat S, Cesur G, Koras O, Degirmenci T. Does Access Location Influence Outcome of Percutaneous Nephrolithotomy in Staghorn Stones?. *J Coll Physicians Surg Pak* 2021; **31(09)**:1075-1080.

INTRODUCTION

Staghorn stones are branched and usually infected stones that occupy a large part of the collecting system.¹ Failure to remove staghorn stones may result in the destruction of the renal parenchyma, leading to the complete loss of kidney function and development of sepsis. Guidelines recommend percutaneous nephrolithotomy (PCNL) as the gold standard treatment for kidney stones greater than 2cm.² The overall success in patients undergoing PCNL has been reported to reach 96.1%.³

However, since more than one percutaneous access may be required to extract all stone pieces in staghorn stones, it is very difficult to reach stone-free status in a single access in these patients.⁴ A previous study reported that the stone-free rate in staghorn stones was decreased to 56.9%.⁵ In addition to these low stone-free rates after PCNL, staghorn stones also have high complication rates. In a prospective randomized study, the intraoperative complication rate of PCNL applied in the treatment of staghorn stones was found to be 16.3%, and the major postoperative complication rate was determined as 18.6%.⁶

The access site in PCNL depends on various factors, such as surgeon preference, renal anatomy, stone location, and total stone burden. In studies, generally accepted view is that calyces other than the lower calyx allow for easier dilatation through the placement of the guide into the ureter from the ureteropelvic junction, resulting in less manipulative trauma and providing a higher stone-free rate. However, the possibility of intercostal access increases in upper calyceal access, which increases the possibility of bleeding due to thoracic complications, including pneumothorax and hydrothorax.⁷⁻¹⁰ On the

Correspondence to: Dr. Murat Sahan, Department of Urology, HSU Izmir Bozyaka Training and Research Hospital, Izmir, Turkey
E-mail: muratsahan87@hotmail.com

Received: June 11, 2021; Revised: July 29, 2021;

Accepted: August 26, 2021

DOI: <https://doi.org/10.29271/jcpsp.2021.09.1075>

other hand, in lower calyceal access, there is a higher risk of colonic injury.¹¹ As stated by Miller *et al.*, access site should facilitate the use of rigid instruments, increase the stone-free rate, and reduce the complication rate to the lowest levels.¹²

The aim of this study was to compare the outcomes of lower and middle calyceal accesses in single-access PCNL performed in a challenging patient group, *i.e.* staghorn stones.

METHODOLOGY

This study was a retrospective analysis of patients who underwent PCNL for staghorn stones at the Izmir Bozyaka Training and Research Hospital from April 2012 to January 2019. The study was approved by the Hospital Ethical Board (Meeting/Decision No.2021/71). All the data of the patients were obtained from electronic medical records.

Patients with congenital kidney anomalies, coagulopathy, skeletal deformity, and solitary kidney, those without staghorn stones, and cases in which multiple accesses used, were excluded from the study. The patients were grouped as lower and middle calyx access: Group-1 (middle calyceal access) and Group-2 (lower calyceal access). Demographic and perioperative data were compared between these two groups.

All patients were evaluated preoperatively using standard non-contrast abdominal computed tomography (CT). The patients' data were recorded by the surgeon. Preoperatively, all patients signed an informed consent form; and were evaluated in terms of stone-free status and complication. Stone-free status was accepted as the presence of stones < 4 mm or no stone detected on follow-up. CT undertaken in the first month after surgery. Postoperative complications were analysed according to the Clavien scoring system.¹³

Stone burden was calculated in square millimeters in all patients as follows: width x length x π x 1/4, where π was taken as 3.14.¹⁴ For staghorn stones, this calculation was performed separately for each calyceal location, and the sum of all calculations were accepted as results. All the PCNL operations were performed by experienced urologists.

The operation began with the insertion of a 5 or 6 F ureteric catheter into the renal pelvis in the lithotomy position under general anesthesia. Subsequently, the kidney was accessed from the lower or middle calyx group with an 18-gauge needle in the prone position under fluoroscopy, depending on the location of the stone and the anatomy of the kidney. The site of entry was dilated up to 30 Fr with Amplatz dilators, and a rigid nephroscope was used to enter the pelvicalyceal system. The stones were fragmented using a pneumatic lithotripter (Elmed; Vibrolith; Ankara/Turkey). Single-access PCNL was performed in all patients. A 14 F nephrostomy catheter was placed postoperative in all patients, and antegrade pyelography was performed.

Categorical data were given as numbers and percentages of columns. The Shapiro-Wilk test was used to examine the conformity of the continuous variables to the parametric distribution. Continuous data were presented as mean value and standard

deviation (SD); and median value and inter-quartile range according to whether they fit the normal distribution or not. In the comparison of continuous data, either Mann-Whitney U-test or Independent-samples t-test was used as appropriate. Pearson's Chi-square and Fisher's Exact tests were used for inter-group comparison of categorical variables. Univariate and multivariate binary logistic regression analyses were performed to analyse possible predictive factors associated with stone-free status. A p-value of <0.05 was considered statistically significant. SPSS software (version 23.0; IBM Corporation, Armonk, NY, USA) was used for the statistical analyses.

RESULTS

There were 249 patients in the study, 108 were in middle calyceal access group and 141 were in lower calyceal access group. The median age of the patients was 49 (37-60) and 50 (38-57.5) years in Group-1 and Group-2, respectively ($p=0.856$). The number of patients with comorbidities was similar in both groups ($p=0.267$). Stone burden and density were also similar in both the groups ($p=0.876$ and $p=0.352$, respectively, Table I).

Table I: Patients' characteristics between the groups.

	Group 1 (n:108)	Group 2 (n:141)	p-value
Age, (years)	49 (37-60)	50 (38-57.5)	0.856
Gender; male/female, n/n	69/39	99/42	0.291*
BMI, kg/m ²	26.1 (22.6-29.4)	27.4 (23.6-30.4)	0.117
Metabolic Syndrome, n(%)			0.714*
-Absent	103 (95.4)	133 (94.3)	
-Present	5 (4.6)	8 (5.7)	
Comorbidity			0.267*
-Diabetes mellitus	4 (3.7)	15 (10.6)	
-Hypertension	7 (6.5)	12 (8.5)	
-Coronary artery disease	9 (8.3)	13 (9.2)	
- ≥ 2	4 (3.7)	6 (4.3)	
Side; left/right, n/n	55/53	80/61	0.362*
Stone burden, mm ²	765 (524-1322)	777 (490-1445)	0.876
Stone density, HU	1120 (900-1200)	1100 (800-1300)	0.352
Parenchymal thickness, mm	13.6 $\pm$ 4.5	15.0 $\pm$ 4.6	0.060
Previous Ipsilateral Surgery, n(%)			0.089*
-Absent	68 (63.0)	103 (73.0)	
-Present	40 (37.0)	38 (27.0)	
Preoperative Hgb, g/dL	13.9 $\pm$ 1.8	13.9 $\pm$ 1.7	0.345
Preoperative GFR, ml/min	90.2 $\pm$ 30.0	90.9 $\pm$ 29.7	0.868

BMI: Body mass index; HU: Hounsfield unit; GFR: Glomerular filtration rate.

*Pearson Chi-square.

Regarding the perioperative and postoperative outcomes, operative, nephroscopy and fluoroscopy times, and hospitalisation time were similar in both the groups ($p=0.343$, $p=0.330$, $p=0.884$, and $p=0.055$, respectively). The number of cases in which intercostal access was used was significantly higher in middle calyceal access group ($p<0.001$). The median dropping hemoglobin was 1.65 (0.80-2.60) g/dL in Group-1 and 1.90

(1.05-2.70) g/dL in Group-2 ($p=0.322$). The number of patients requiring transfusion replacement was similar in both the groups ($p=0.346$, Table II).

Table II: Perioperative ve postoperative outcomes.

	Group 1 (n:108)	Group 2 (n:141)	p-value
Access level, n(%)			
-11 th -12 th intercostal	62 (57.4)	29 (20.6)	<0.001*
-Subcostal	46 (42.6)	112 (79.4)	
Operative time, (min)	120 (90-150)	120 (90-145)	0.343
Nephroscopy time, (min)	50 (40-90)	50 (40-70)	0.330
Fluoroscopy time, (sec)	65 (45-92)	68 (40-98.5)	0.884
Hospitalization time, (day)	4 (3-5)	3 (2-4.5)	0.055
Hgb drop, (g/dL)	1.65 (0.80-2.60)	1.90 (1.05-2.70)	0.322
GFR change, ml/min	-11.0 (-18.0 to -2.41)	-9.4 (-17.2 to 0)	0.278
Transfusion rates, n(%)	14 (13.0)	13 (9.2)	0.346*
Stone free rates, n(%)	50 (46.3)	93 (66.0)	0.002*
Clavien-Dindo classification, n(%)			0.205*
-No complication	66 (61.1)	99 (70.2)	
-Clavien I-II	27 (25.0)	31 (22.0)	
-Clavien III-V	15 (13.9)	11 (7.8)	

Hgb: Hemoglobin; GFR: Glomerular filtration rate. * Pearson Chi-square

Table III: Classification of complications according to Clavien scoring system.

Complications	Group 1 (n:42)	Group 2 (n:42)	p value
Grade I, n (%)	3 (7.1)	9 (26.2)	0.061*
-Pain	-	4	
-Fever	1	5	
-Bleeding not requiring blood transfusion	2		0.661*
Grade II, n (%)	24 (57.1)	22 (47.6)	
-Bleeding requiring blood transfusion	9	13	
-Postoperative fever (>38°C) managed with antibiotics in the ward	15	9	
Grade III	14 (33.3)	11 (26.2)	0.474*
Grade III A, n(%)			
-Urine leakage managed by ureteric stenting without general anesthesia	7	7	
-Perinephric abscess managed by percutaneous drainage	2	1	0.999^
-Hydrothorax	2	-	
-Renal hemorrhage requiring angioembolization	-	1	
Grade III B, n(%)			0.999^
-Urine leakage managed by ureteric stenting with general anesthesia	3	2	
Grade IV	1 (2.4)	0	
Grade IV B, n (%)			0.999^
-Sepsis	1	-	

* Pearson Chi-square; ^ Fisher's Exact Test.

Stone-free status was detected in 50 (46.3%) patients in lower calyceal access group and 93 (66.0%) patients in middle calyceal access group ($p=0.002$). Patients who had clinically significant residual stones after single-access PCNL, were undergone extracorporeal shock wave lithotripsy (19 patients), retrograde intrarenal surgery (32 patients), and second look-PCNL (8 patients). Concerning the overall complication rates, complications were detected in 42 (38.9%) patients in Group-1 and 42 (29.8%) patients in Group-2 ($p=0.132$, Table II).

When the authors examined complication subgroups formed according to the Clavien scoring system, they observed that the two groups had a similar distribution of complication grades (Table III).

In the univariate analysis, stone density (OR: 0.999, $p=0.025$), stone burden (OR: 1.000 $p=0.038$), and lower calyceal access (OR: 2.247, $p=0.002$) were found to be associated with stone-free status. The multivariate analysis conducted with stone burden (OR: 1.000 $p=0.045$), and lower calyceal access (OR: 2.289 $p=0.002$) were independent risk factors for success.

DISCUSSION

Despite advances in instrumentation and technology, staghorn stones are still difficult to manage. PCNL is a preferred treatment method with a high success and low major complication rate in the treatment of complex kidney stones.¹⁵ However, there were some studies which showed that the success rate in staghorn kidney stones was decreased to 56.9%.⁵ Although some authors suggest the superiority of access through certain calyces for routine PCNL, there is an ongoing debate concerning which renal calyceal access achieves higher success in staghorn stones. In a retrospective evaluation of 153 PCNL procedures with ultrasound-guided access, the authors reported that the middle calyceal access had a shorter operative time (41 minutes) and higher stone-free rate (98%) compared to the lower (50 min and 84%, respectively) and upper (46 min and 93%, respectively) calyceal access groups, but they observed no significant difference between the groups in relation to complications.¹⁶ In the CROES PCNL Global study, upper and lower pole accesses were compared in 4,494 patients that underwent single-access PCNL, and it was observed that the upper pole access group had a extended operative time and lower stone-free rate (77 versus 82%), higher overall complication rate (23.5 versus 16%), and longer hospital stay.¹⁷ To the best of authors' knowledge, this study was the first study to assess the effect of middle and lower calyceal accesses on the outcomes of staghorn kidney stone treatment.

In this study, which aimed to compare the safety and efficiency of middle and lower pole accesses, only those cases were evaluated in which the surgeon considered only single access was required; and excluded those involving multiple entries. In a study by Verma *et al.* conducted with 110 PCNL cases with single and multiple accesses, it was observed that patients who required only one access site for stone removal had a larger infundibular area, greater intercalyceal angle, and smaller pelvicalyceal surface area.¹⁸ Although these parameters were not clearly measured in this study, such evaluation was made by retrograde pyelography performed in each case. When evaluating the access site in PCNL, it is important to know that kidney anatomy may differ between patients.

Although previous studies have reported higher stone-free rates (78-93%) after PCNL in staghorn kidney stones,^{4,19} in the current study including staghorn stones, the success rate was

66.0 % in the lower calyceal access group and 46.3% in the middle calyceal access group ($p=0.002$). This lower rate of success can be explained by the exclusion of use of a single access in each patient, lithotripsy being performed with only a pneumatic lithotripter, and technical limitations caused by not using a flexible nephroscope. Previous studies have shown that middle calyceal access facilitates entry into many points of the pelvicalyceal system by providing greater maneuverability and is more advantageous than lower calyceal access, especially in staghorn and complex stones.¹⁶ In contrast, Blum *et al.* found the stone-free rate to be 77.3% in patients with a single lower calyceal access in staghorn kidney stones, while they found a stone-free rate of 68.2% in the single upper calyceal access group.²⁰ Similarly, in this study, the stone-free rate was significantly higher in the lower calyceal access group. Choosing an appropriate access site is an important factor influencing success, but it is not possible to define it precisely since there are many other factors affecting success, such as different renal anatomy, stone size, density and type, and technical equipment used (flexible nephroscope, etc.). In present study, the authors found that stone burden, and lower calyceal access were independent risk factors for success. In a series of 498 patients, Binbay *et al.* examined the anatomical variations of the kidney and reported a positive correlation between a lower pelvicalyceal surface area and stone configuration and a high stone-free rate.²¹

In this study, in which only staghorn stones were included, there was no significant difference in the stone size of the two groups, which also shows the comparability of the groups in term of stone burden. In the present study, length of hospital stay, operative, fluoroscopy nephroscopy time, hemoglobin drop, and blood transfusion rate were found to be similar in both the groups. The access site being the lower or middle calyx has no effect on these parameters in complex stones. However, Caglayan *et al.* reported that factors such as renal parenchymal damage, forced tract dilatation, sharp renal axis angle, and increased torque due to limited manipulation in the lower calyceal access group might be associated with greater blood loss, increased number of residual stones, and longer operative time.²² In another study, it was shown that the renal pelvis and upper calyx could be accessed more easily using the posterior lower calyceal route and less torque would result in less damage to the renal parenchyma and infundibulum. It was also shown that the more medial approach can facilitate laterally located anterior lower calyceal access, and the use of a proper access angle into the infundibulum could provide better access to the rest of the collector system, including the interpolar calyces. In addition, an access angle that is less perpendicular to the collector system was suggested to facilitate the gravitational passage of materials through the sheath.²¹ Karve *et al.* reported that hemoglobin drop was 0.67 g/dL in patients with a middle calyceal access and 1.36 g/dL in those with a lower calyceal access ($p<0.001$). The authors reported that transfusion rates were similar in both groups²³. Similarly, in another study, while hemoglobin drop was significantly higher in the

lower calyceal access group, there was no statistically significant difference in transfusion rates of the groups.¹⁶

The goal of upper urinary tract stone treatment methods is to provide maximum stone-free rates with minimum morbidity. PCNL is considered as an effective and safe treatment option for large kidney stones. Despite the proven efficacy of the PCNL, complication rates reaching 83% have been reported in the literature, including bleeding requiring transfusion (7%), organ damage (0.4%), and infectious events (up to 33%).²⁴ In this study, the two groups had similar rates of major (13.9% in Group-1 and 7.8% in Group-2) and minor (25.0 and 22.0%, respectively) complications. The overall complication rate was 38.9% in middle calyceal access group and 29.8% in lower calyceal access group; and there was no statistically significant difference between them. When the complication subgroups were examined according to the Clavien scoring system, it was seen that postoperative fever requiring antibiotics was more common than the other complications. This may be related to staghorn stones, generally being infected stones, and the presence of residual stones causing infectious events. Choosing an appropriate access site during PCNL plays a critical role in the success of the operation. Upper calyceal access provides direct entry into all renal calyces and the pelvis; however, the increased possibility of thoracic complications remains a major concern for urologists. Soares *et al.* showed that supracostal upper calyceal access was associated with a higher rate of thoracic complications.²⁵ In this study, it was observed that hydrothorax developed in two cases in which supracostal middle calyceal access had been used. In light of this information, lower calyx of entry during the PCNL procedure should be preferred in staghorn kidney stones due to high success and similar complication rates.

There are some limitations to the generalisation of this study. First, it had a retrospective design and a limited number of patients. Second, there was no long-term results of surgical complications. Finally, there is a need for prospective studies with a larger series of patients with staghorn stones, focusing on the access site.

CONCLUSION

This study showed that, in PCNL applied to staghorn kidney stones using a single access, lower calyceal access was superior to middle calyceal access in relation to the stone-free rates. The selection of an appropriate access site in the treatment of staghorn kidney stones may vary according to patient factors and surgeon experience. The results of this study should be confirmed by further studies evaluating the efficacy of different calyceal access sites in staghorn kidney stones.

ETHICAL APPROVAL:

This study follows the principles of the Declaration of Helsinki; and it was approved by the Ethics Committee of the Turkish Ministry of Health, Izmir Bozyaka Training and Research Hospital (Meeting/Decision No. 2021/71).

PATIENTS' CONSENT:

Informed consent was not obtained due to the retrospective nature of the study design.

CONFLICT OF INTEREST:

The authors declared no conflict of interest.

AUTHORS' CONTRIBUTION:

SY: Study conception and design, analysis and interpretation of data, wrote the paper.

MS: Study conception and design, wrote the paper.

SP: Acquisition of data, analysis and interpretation of data.

OK: Acquisition of data, critical revision.

GC: Analysis and interpretation of data, statistical analysis.

TD: Analysis and interpretation of data, critical revision.

REFERENCES

- Ganpule AP, Desai M. Management of the staghorn calculus: Multiple-tract versus single-tract percutaneous nephrolithotomy. *Curr Opin Urol* 2008; **18(2)**:220-3. doi: 10.1097/MOU.0b013e3282f3e6e4.
- Türk C, Petřík A, Sarica K, Seitz C, Skolarikos A, Straub M, et al. eau guidelines on interventional treatment for urolithiasis. *Eur Urol* 2016; **69(3)**:475-82.
- Wollin DA, Preminger GM. Percutaneous nephrolithotomy: Complications and how to deal with them. *Urolithiasis* 2018; **46(1)**:87-97. doi: 10.1007/s00240-017-1022-x.
- Desai M, Jain P, Ganpule A, Sabnis R, Patel S, Shrivastav P. Developments in technique and technology: The effect on the results of percutaneous nephrolithotomy for staghorn calculi. *BJU Int* 2009; **104(4)**:542-8. doi: 10.1111/j.1464-410X.2009.08472.x.
- Desai M, De Lisa A, Turna B, Rioja J, Walfridsson H, D'Addessi A. The clinical research office of the endourological society percutaneous nephrolithotomy global study: Staghorn versus non staghorn stones. *J Endourol* 2011; **25(8)**:1263-8. doi: 10.1089/end.2011.0055.
- Al-Kohlany KM, Shokeir AA, Mosbah A, Mohsen T, Shoma AM, Eraky I, et al. Treatment of complete staghorn stones: A prospective randomized comparison of open surgery versus percutaneous nephrolithotomy. *J Urol* 2005; **173(2)**:469-73. doi: 10.1097/01.ju.0000150519.49495.88.
- Preminger GM, Assimos DG, Lingeman JE, Nakada SY, Pearle MS, Wolf JS Jr. AUA (nephrolithiasis guideline panel). Chapter 1: AUA guideline on management of staghorn calculi: diagnosis and treatment recommendations. *J Urol* 2005; **173(6)**:1991-2000. doi: 10.1097/01.ju.0000161171.67806.2a.
- Shah HN, Hegde SS, Shah JN, Bansal MB. Safety and efficacy of supracostal access in tubeless percutaneous nephrolithotomy. *J Endourol* 2006; **20(12)**:1016-21. doi: 10.1089/end.2006.20.1016.
- Sukumar S, Nair B, Ginil KP, Sanjeevan KV, Sanjay BH. Supracostal access for percutaneous nephrolithotomy: Less morbid, more effective. *Int Urol Nephrol* 2008; **40(2)**:263-7. doi: 10.1007/s11255-007-9270-2.
- Munver R, Delvecchio FC, Newman GE, Preminger GM. Critical analysis of supracostal access for percutaneous renal surgery. *J Urol* 2001; **166**:1242-6.
- Kyriazis I, Panagopoulos V, Kallidonis P, Özsoy M, Vasilas M, Liatsikos E. Complications in percutaneous nephrolithotomy. *World J Urol* 2015; **33**:1069-77.
- Miller NL, Matlaga BR, Lingeman JE. Techniques for fluoroscopic percutaneous Renal access. *J Urol* 2007; **178(1)**:15-23. doi: 10.1016/j.juro.2007.03.014.
- Dindo D, Demartines N, Clavien PA. Classification of surgical complications: A new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg* 2004; **240(2)**:205-13. doi: 10.1097/01.sla.0000133083.54934.ae.
- Tiselius HG, Andersson A. Stone burden in an average Swedish population of stone formers requiring active Stone removal: How can the stone size be estimated in the clinical routine? *Eur Urol* 2003; **43(3)**:275-81. doi: 10.1016/s0302-2838(03)00006-x.
- de la Rosette J, Assimos D, Desai M, Gutierrez J, Lingeman J, Scarpa R, et al. Croes PCNL study group. The clinical research office of the endourological society percutaneous nephrolithotomy global study: Indications, complications, and outcomes in 5803 patients. *J Endourol* 2011; **25(1)**:11-7. doi: 10.1089/end.2010.0424.
- Song Y, Jin W, Hua S, Fei X. Middle calyx access is better for single renal pelvic stone in ultrasound-guided percutaneous nephrolithotomy. *Urolithiasis* 2016; **44(5)**:459-63. doi: 10.1007/s00240-016-0866-9.
- Tefekli A, Esen T, Olbert PJ, Tolley D, Nadler RB, Sun YH, et al. Croes PCNL study group. Isolated upper pole access in percutaneous nephrolithotomy: A large-scale analysis from the croes percutaneous nephrolithotomy global study. *J Urol* 2013; **189(2)**:568-73. doi: 10.1016/j.juro.2012.09.035.
- Verma A, Tomar V, Yadav S. Complex multiple renal calculi: Stone distribution, pelvicalyceal anatomy and site of puncture as predictors of PCNL outcome. *Springer Plus* 2016; **5(1)**:1356. doi: 10.1186/s40064-016-3017-4.
- Soucy F, Ko R, Duvdevani M, Nott L, Denstedt JD, Razvi H. Percutaneous nephrolithotomy for staghorn calculi: A single center's experience over 15 years. *J Endourol* 2009; **23(10)**:1669-73. doi: 10.1089/end.2009.1534.
- Blum KA, Parkhomenko E, Thai J, Tran T, Gupta M. A contemporary lower pole approach for complete staghorn calculi: Outcomes and efficacy. *World J Urol* 2018; **36(9)**:1461-7. doi: 10.1007/s00345-018-2284-9.
- Binbay M, Akman T, Ozgor F, Yazici O, Sari E, Erbin A, et al. Does pelvicalyceal system anatomy affect success of percutaneous nephrolithotomy? *Urology* 2011; **78(4)**:733-7. doi: 10.1016/j.urology.2011.03.058.
- Caglayan V, Onen E, Avci S, Kilic M, Sambel M, Oner S. Percutaneous nephrolithotomy via a middle calyx access is effective in the treatment of lower pole kidney stones: A single-center study. *Urol Int* 2020; **104(9-10)**:741-5. doi: 10.1159/000505081.
- Falahatkar S, Kazemnezhad E, Moghaddam KG, Kazemzadeh M, Asadollahzade A, FarzanA, et al. Middle calyx access in complete supine percutaneous nephrolithotomy. *Can Urol Assoc J* 2013; **7(5-6)**:E306-10. doi: 10.5489/cuaj.11307.

24. Michel MS, Trojan L, Rassweiler JJ. Complications in percutaneous nephrolithotomy. *Eur Urol* 2007; **51**:899-906.
25. Soares RMO, Zhu A, Talati V, Nadler RB. Upper pole access for prone percutaneous nephrolithotomy: Advantage or risk? *Urology* 2019; **134**:66-71. doi: 10.1016/j.urology.2019.08.031.

• • • • •