## LETTER TO THE EDITOR

# Collapsibility and Distensibility Indices of Critically III Patients

Sir,

I have read with great interest the recent article, "Inferior Vena Cava Collapsibility Index and Central Venous Pressure for Fluid Assessment in the Critically III Patient" by Dodhy AA.<sup>1</sup> This article is a well-designed cross-sectional study for the detection of the collapsibility index of inferior vena cava (IVC) in critically III patients. Also, this topic is very important for the emergency departments and intensive care units (ICUs). The use of point-of-care ultrasound for the critically ill patients by emergency physicians and intensivists has become widespread.

In the study of Dodhy, the researcher evaluated the diameter and collapsibility index of IVC in 126 patients and compared and correlated it with central venous pressure (CVP) in both mechanically ventilated and spontaneous breathing groups. There are some points I would like to point out about the study. The collapsibility index is a volume indicator that has been used safely in patients with spontaneous breathing for several years.<sup>2,3</sup> Positive-pressure ventilation elevates the pleural and right atrial pressures. Also, the venous return to the heart reduces by increasing intrathoracic pressure during inspiration. The IVC diameter increases during inspiration and contracts during expiration in a mechanically ventilated patient, unlike in spontaneous breathing.<sup>4</sup> Since the researcher evaluated only IVC diameter and collapsibility index, I think that the distensibility index should be added to the study. Since the majority of the patients were mechanically ventilated, the distensibility index might give a different correlation coefficient.

Nonetheless, I think it is a very important article about the point-of-care ultrasound performed in the ICU. I think that it should be recommended to use the distensibility index as a non-invasive method in mechanically ventilated patients due to physiological changes.

### **COMPETING INTEREST:**

The author declared no competing interest.

### **AUTHOR'S CONTRIBUTION:**

MEC: Conceptualisation, writing, original draft, review and editing.

### REFERENCES

1. Dodhy AA. Inferior vena cava collapsibility index and central venous pressure for fluid assessment in the

critically III patient. *J Coll Physicians Surg Pak* 2021; **31(11)**: 1273-7. doi: 10.29271/jcpsp.2021.11.1273.

- Barbier C, Loubières Y, Schmit C, Hayon J, Ricôme J-L, Jardin F, *et al.* Respiratory changes in inferior vena cava diameter are helpful in predicting fluid responsiveness in ventilated septic patients. *Intensive Care Med* 2004; **30(9)**:1740-6. doi: 10.1007/s00134-004-2259-8.
- Gui J, Yang Z, Ou B, Xu A, Yang F, Chen Q, *et al.* Is the collapsibility index of the inferior vena cava an accurate predictor for the early detection of intravascular volume change? *Shock* 2018; **49(1)**:29-32. doi: 10.1097/SHK. 00000000000932.
- 4. Tan HL, Wijeweera O, Onigkeit J. Inferior vena cava guided fluid resuscitation fact or fiction? *Trends Anaesthesia Critical Care* 2015; **5(2)**:70-5.

#### Mustafa Emin Canakci

Department of Emergency Medicine, Eskisehir Osmangazi University, Eskisehir, Turkey

Correspondence to: Dr. Mustafa Emin Canakci, Department of Emergency Medicine, Eskisehir Osmangazi University, Eskisehir, Turkey E-mail: mustafaeminc@gmail.com

Received: December 06, 2021; Revised: December 09, 2021; Accepted: December 25, 2021 DOI: https://doi.org/10.29271/jcpsp.2022.07.953

### **AUTHOR'S REPLY**

It is a good suggestion by the reader that the distensibility index be added in the study for mechanically ventilated patients and the results may differ. However, in the present study, the aim was to determine the correlation of sonographic evaluation of IVC diameter and its collapsibility index with CVP in both spontaneously breathing and mechanically ventilated patients and thus to evaluate the intravascular volume status of critically ill patients. Abdelwahab and El-Wahab, in their study, also aimed to detect the correlation between IVC diameter and collapsibility index with CVP in both spontaneously breathing and mechanically ventilated patients.<sup>1</sup>Karacabey *et al.* evaluated the strong correlation between CVP and IVC diameters and collapsibility in intubated patients as assessed by ultrasonography.<sup>2</sup> The suggestion of the reader can be incorporated in future research papers.

### REFERENCES

1. Abdelwahab HW, El-Wahab RA. Correlation of bedside ultrasonographic measurement of IVC diameter and caval index with CVP in spontaneously breathing and mechanically ventilated patients. *J Emerg Med Critical Care* 2017; **3(1)**:2.  Karacabey S, Sanri E, Guneysel O. A non-invasive method for assessment of intravascular fluid status: Inferior vena cava diameters and collapsibility index . *Pak J Med Sci* 2016; **32(4)**:836-40. doi.org/10.12669/ pjms.324.10290.

.....

Afia Arshed Dodhy

Department of Anaesthesia and ICU, Lahore General Hospital, PGMI / LGH / AMC, Lahore, Pakistan

.....

Correspondence to: Dr. Afia Arshed Dodhy, Department of Anaesthesia and ICU, Lahore General Hospital, PGMI / LGH / AMC, Lahore, Pakistan E-mail: afiashujaat@yahoo.com

• • • • • • • • • • •