Diagnostic Value of Serum Gastrin and Epidermal Growth Factor to the Gastric Ulcer Complicated with Upper Gastrointestinal Hemorrhage

Ying Li and Yuping Song

Department of Emergency, Beijing Jiangong Hospital, China

ABSTRACT

Objective: To explore the predictive value of serum gastrin (GAS), epidermal growth factor (EGF) levels in gastric ulcer complicated with acute upper gastrointestinal bleeding.

Study Design: A descriptive study.

Place and Duration of Study: Department of Emergency, Beijing Jiangong Hospital, China, from January 2019 to June 2020.

Methodology: One hundred and twenty-five patients with gastric ulcer and acute upper gastrointestinal bleeding were selected as Group A. One hundred and twenty-five patients with gastric ulcer and no upper gastrointestinal bleeding were selected as Group B. Logistic regression analysis was used to analyse the risk factors of gastric ulcer complicated with acute upper gastrointestinal bleeding. The value of serum GAS, EGF in early diagnosis of gastric ulcer with upper gastrointestinal bleeding was evaluated by receiver operating characteristic (ROC) curve.

Results: Univariate analysis showed statistically significant differences between Group A and Group B in taking non-steroidal anti-inflammatory drugs (NSAIDs), *helicobacter pylori* (Hp) infection, serum GAS and EGF (all $p < 0.001$). Logistic regression analysis showed that raised serum GAS and serum EGF were independent risk factors for gastric ulcer and upper gastrointestinal bleeding (both $p < 0.001$). The ROC area of serum EGF to predict gastric ulcer and acute upper gastrointestinal bleeding was 0.810 (95% CI: 0.753-0.867, $p < 0.001$), greater than ROC area of serum GAS. At serum EGF of ≤ 109.95 pg/mL, had the 84.8%, sensitivity to predict gastric ulcer and acute upper gastrointestinal bleeding with specificity of 68.8%.

Conclusion: The predictive value of serum GAS and EGF is high for gastric ulcer complicated with acute upper gastrointestinal bleeding; the predictive value of serum EGF is greater than that of serum GAS.

Key Words: Gastric ulcer, Acute upper gastrointestinal bleeding, Serum, Gastrin (GAS), Epidermal growth factor (EGF), Logistic regression, Receiver operating characteristic (ROC) curve.

How to cite this article: Li Y, Song Y. Diagnostic Value of Serum Gastrin and Epidermal Growth Factor to the Gastric Ulcer Complicated with Upper Gastrointestinal Hemorrhage. *J Coll Physicians Surg Pak* 2020; **30(12)**:1269-1272.

INTRODUCTION

Gastric ulcer mainly refers to tissue damage exceeding mucosal muscular layer, caused by gastric mucosa digestion by gastric digestive juice itself. It is an inflammatory necrotic lesion that occurs between the cardia and the pylorus.¹ Peptic ulcer is a common cause of upper gastrointestinal hemorrhage.² Acute upper gastrointestinal bleeding is a common complication of gastric ulcer. Most of the times, the bleeding volume is 300-500 mL. When the bleeding volume is more than 800 mL, systemic shock and even life-threatening may occur.³

Gastroscopy, also known as upper gastrointestinal endoscopy, is currently the preferred examination method for the etiology and location of upper gastrointestinal hemorrhage in diagnosis. An examination performed within 24-48 hours after hemorrhage is called an emergency gastroscopy. Upper gastrointestinal hemorrhage can be diagnosed by gastroscopy.⁴ Early gastroscopy and accurate diagnosis are the key to reduce mortality. Compared with endoscopy, blood examination has the advantages of low cost and low risk.⁵

Gastrin (GAS) is a peptide hormone synthesised and secreted by gastric antral G cells. After release, it acts on gastric parietal cells to stimulate the secretion of hydrochloric acid through improving the blood circulation, which plays an important role in regulating digestive function.⁶ One study has confirmed that serum GAS level in patients with gastric ulcer has increased.⁷

Epidermal growth factor (EGF) is a kind of endogenous protective substance, which can promote the proliferation, differentiation and migration of epithelial cells. EGF has the repairing and

Correspondence to: Yuping Song, Department of Emergency, Beijing Jiangong Hospital, 100054, China
E-mail: iuf535@163.com

Received: October 10, 2020; Revised: November 15, 2020;
Accepted: November 24, 2020
DOI: <https://doi.org/10.29271/jcpsp.2020.12.1269>

protecting function to gastric mucosa damage.⁸ A study has shown that serum EGF is involved in the occurrence and healing of gastric ulcers.⁹

At present, the application value of serum GAS and EGF in acute upper gastrointestinal bleeding, complicated with gastric ulcer, is not yet clear. In view of this, the purpose of this study was to explore the predictive value of serum GAS and EGF levels in gastric ulcer complicated with acute upper gastrointestinal bleeding.

METHODOLOGY

This descriptive study was approved by The Ethics Committee of Beijing Jiangong Hospital, China. A total of 125 patients with gastric ulcer and acute upper gastrointestinal bleeding admitted to the hospital from January 2019 to June 2020, were selected as Group A. One hundred and twenty-five patients with gastric ulcer and no upper gastrointestinal bleeding admitted to the hospital during the same period, were selected as Group B.

Inclusion criteria were that patients with gastric ulcer diagnosed by gastroscopy, without effective treatment before admission, complete medical history information, voluntary acceptance of this trial. Exclusion criteria were that patients aged >80 or <18 years; diagnosed with duodenal ulcer, pyloric duct ulcer and compound ulcer by gastroscopy; received peptic endoscopic hemostasis treatment within two months; combined with severe respiratory, circulation, hepatic or renal dysfunction, upper gastrointestinal bleeding caused by liver cirrhosis, gastric cancer, acute gastric mucosal lesions, etc.; pregnant or breastfeeding women; bad addictions (such as alcohol and drug abuse); mental illness or with impaired consciousness.

Relevant medical information of patients in Group A and Group B were collected including gender, age, whether they have taken NSAIDs, *helicobacter pylori* (Hp) infection, etc.

The urea breath test was used to detect Hp infection. The subjects had to fast for at least 4 hours before detection. After taking off the cap of the bottom air bag, the subjects blew the air bag full and tightened the cap. Then they took a urea (¹⁴C) capsule. After resting for 15 minutes, they blew again with a gas collecting card in the mouth. They could breathe during the blowing process, and they stopped when the gas collecting card changed from orange to yellow. The subjects' gas collection card was detected by ¹⁴C detector, and Hp was positive when the judgment value was equal to or greater than 50.

Upon admission, 3 mL venous blood was drawn from the subjects, and centrifuged to obtain serum. Enzyme-linked immunosorbent assay (ELISA) was used to detect serum EGF levels, and radioimmunoassay was used to determine serum GAS levels.

Data was analysed using SPSS 25.0. Kolmogorov-Smirnov test was used to test the normality of all measurement data. Measurement data such as age, serum GAS, serum EGF were expressed as mean ± SD, and were compared by independent

sample t-test between the two groups. Categorical data were expressed as frequencies and percentages. Categorical data between the two groups were compared by Chi-square test. Logistic regression analysis was used to analyse the risk factors of gastric ulcer complicated with acute upper gastrointestinal bleeding. The value of serum GAS, EGF in early diagnosis of gastric ulcer with upper gastrointestinal bleeding was evaluated by receiver operating characteristic (ROC) curve. A p-value less than 0.05 was regarded as statistically significant.

RESULTS

The results of univariate analysis showed that there were statistically significant differences between Group A and Group B in taking NSAIDs, Hp infection, serum GAS and EGF (all p < 0.001, Table I); however, there was no significant difference in gender, age, smoking and drinking groups (p = 0.692, 0.660, 0.163 and 0.097, respectively, Table I).

Table I: Comparison of related parameters in the two groups.

Index	Group A (n=125)	Group B (n=125)	p-value
Gender [male, n (%)]	79 (63.20)	82 (65.60)	0.692
Age (years)	58.08±9.48	57.56±9.20	0.660
Smoking [n (%)]	73 (58.40)	62 (49.60)	0.163
Drinking [n (%)]	77 (61.60)	64 (51.20)	0.097
Taking NSAIDs [n (%)]	106 (84.80)	70 (56.00)	<0.001
Hp infection [n (%)]	114 (91.20)	88 (70.40)	<0.001
Serum GAS (μg/L)	94.33±15.32	123.67±19.78	<0.001
Serum EGF (pg/mL)	100.75±16.41	138.50±22.15	<0.001

Logistic regression analysis was performed on factors such as taking NSAIDs (yes=1, no=0), Hp infection (yes=1, no=0), serum GAS (actual test value), serum EGF (actual test value), and the results showed that serum GAS and serum EGF were independent risk factors for gastric ulcer and upper gastrointestinal bleeding (both p < 0.001, Table II).

Table II: Multivariate logistic regression analysis of gastric ulcer and upper gastrointestinal bleeding.

Risk factor	B	S.E.	p-value	Exp (B)
Taking NSAIDs	-0.968	1.013	0.339	0.380
Hp infection	-0.908	0.980	0.354	0.403
Serum GAS	2.124	0.328	<0.001	8.361
Serum EGF	-1.864	0.283	<0.001	0.155

The ROC area of serum GAS to predict gastric ulcer and acute upper gastrointestinal bleeding was 0.774 (95% CI: 0.714 - 0.835, p < 0.001). When serum GAS was ≤ 98.09 μg/L, its sensitivity to predict gastric ulcer and acute upper gastrointestinal bleeding was 84.8%, and the specificity was 56%.

The ROC area of serum EGF to predict gastric ulcer and acute upper gastrointestinal bleeding was 0.810 (95% CI: 0.753 - 0.867, p < 0.001). When serum EGF was ≤ 109.95 pg/mL, its sensitivity to predict gastric ulcer and acute upper gastrointestinal bleeding was 84.8%, and the specificity was 68.8%, as shown in Figure 1.

The ROC area of serum EGF to predict gastric ulcer and acute

upper gastrointestinal bleeding was 0.810 (95% CI: 0.753 - 0.867, $p < 0.001$), higher than ROC area of serum GAS.

Figure 1: ROC curves of serum GAS and EGF to predict gastric ulcer and acute upper gastrointestinal bleeding.

DISCUSSION

Hp infection and taking NSAIDs are two important pathogenic factors of peptic ulcer.^{10,11} Numerous studies have confirmed that Hp and NSAIDs are two independent gastric mucosal damaging factors, and can cause gastric mucosal damage through different mechanisms.^{12,13}

NSAIDs mainly inhibit the synthesis of prostaglandins in gastric mucosa, thus inhibiting the secretion of mucus and bicarbonate in gastric epithelial cells, reducing the blood flow of gastric mucosa and the renewal rate of gastric epithelium, stimulating the adhesion of neutrophils in blood vessel wall, and migrating out of blood vessels to infiltrate locally, causing gastric mucosa damage.¹⁴ However, the role of Hp infection in ulcer with hemorrhage has not yet been completely defined. A study has found that Hp infection increases the risk of bleeding peptic ulcer disease.¹⁵

The results showed that there was a statistically significant difference between Group A and Group B in taking NSAIDs and Hp infection in univariate analysis. Multivariate logistic regression analysis showed that Hp infection and recent NSAIDs taken history were not independent risk factors for gastric ulcer combined with upper gastrointestinal bleeding. However, Chen *et al.* found that Hp infections were independent ulcer risk factors using multiple logistic regression analysis.¹⁶ Previous studies have concluded that among patients with peptic ulcer and upper gastrointestinal bleeding, the Hp positive infection rate is significantly higher than that of patients without upper gastrointestinal bleeding.¹⁷

According to a study, serum GAS value was closely related to the disease process of peptic ulcer and was of prognostic value.¹⁸ Research showed that except for Zollinger-Ellison syndrome, a fasting serum GAS level could not be used as a screening test for

peptic ulcer disease in children.¹⁹ The results of this study showed that the difference in serum GAS between Group A and Group B was statistically significant. Multivariate logistic regression analysis showed that serum GAS was an independent risk factor for gastric ulcer and acute upper gastrointestinal bleeding. ROC curve analysis showed that the area under the curve of serum GAS to predict gastric ulcer and acute upper gastrointestinal bleeding was 0.774 (95% CI: 0.714-0.835, $p < 0.001$). When serum GAS was $\leq 98.09 \mu\text{g/L}$, its sensitivity to predict gastric ulcer and acute upper gastrointestinal bleeding was 84.8%, and the specificity was 56%.

One study showed that EGF was important for gastric ulcer healing.²⁰ Serum EGF levels may be of closely relation between gut hormones and peptic ulcer.²¹ The results of this study showed that there was a statistically significant difference in serum EGF between Group A and Group B. Multivariate logistic regression analysis showed that serum EGF was an independent risk factor for gastric ulcer and acute upper gastrointestinal bleeding.

ROC curve analysis showed that the area under the curve of serum EGF to predict gastric ulcer and acute upper gastrointestinal bleeding was 0.810 (95% CI: 0.753-0.867, $p < 0.001$). When serum EGF was $\leq 109.95 \text{ pg/mL}$, its sensitivity to predict gastric ulcer and acute upper gastrointestinal bleeding was 84.8%, and the specificity was 68.8%. The area under the curve of serum EGF to predict gastric ulcer complicated with upper gastrointestinal hemorrhage was larger than that of serum GAS, and its predictive value was greater.

CONCLUSION

Serum GAS and EGF are independent risk factors for gastric ulcer complicated with acute upper gastrointestinal bleeding. The predictive value of serum GAS and EGF is high for gastric ulcer complicated with acute upper gastrointestinal bleeding, and the predictive value of serum EGF is greater than that of serum GAS.

ETHICAL APPROVAL:

This study was conducted with the approval from the Ethics Committee of the Beijing Jiangong Hospital, China.

PATIENTS' CONSENT:

All patients signed a document of informed consent.

CONFLICT OF INTEREST:

The authors have no conflicts of interests to declare.

AUTHORS' CONTRIBUTION:

YL: Interpretation of data for the work, drafting the work.

YS: Revising it critically for important intellectual content, final approval of the version to be published.

REFERENCES

1. Matsueda K, Kawano S, Okada H. Primary localised amyloidosis of the stomach mimicking healing gastric ulcer. *Gastrointest Endosc* 2020; **91**(4):947-8. doi: 10.1016/j.gie.

- 2019.12.004.
2. Ardalani H, Hadipanah A, Amirhossein S. Medicinal plants in the treatment of peptic ulcer disease: A review. *Mini Rev Med Chemis* 2020; **20(8)**:662-702. doi: 10.2174/1389557520666191227151939.
 3. Hreinsson JP, Kalaitzakis E, Gudmundsson S, Björnsson ES. Upper gastrointestinal bleeding: incidence, etiology and outcomes in a population-based setting. *Scand J Gastroenterol* 2013; **48(4)**:439-7. doi: 10.3109/00365521.2012. 763174.
 4. Laursen SB, Gralnek IM, Stanley AJ. Raising the threshold for hospital admission and endoscopy in upper gastrointestinal bleeding during the COVID-19 pandemic. *Endoscopy* 2020; **52(10)**:930-1. doi: 10.1055/a-1202-1374.
 5. Chiang CH, Jeng JE, Wang WM, Jheng BH. A comparative study of three fecal occult blood tests in upper gastrointestinal bleeding. *Kaohsiung J Med Sci* 2006; **22(5)**:223-8. doi: 10.1016/S1607-551X(09)70240-2.
 6. Han H, Sun X, Zheng X, Xiao J. Effects of somatostatin combined with erythromycin on gastrointestinal function recovery, gastrointestinal hormone levels and immune indexes in patients after gastrointestinal surgery. *Acta Microscopica* 2020; **29 (3)**:1454-60.
 7. Kang J, Kang LL, Yu H, Tao C, Chen LM. Levels and significance of serum peptide hormone and monoamine neurotransmitter in patients with gastric ulcer. *Progress Modern Biomedicine* 2016; **16(26)**:5131-4.
 8. Chen W, Wu D, Jin Y, Li Q, Liu Y, Qiao X, et al. Pre-protective effect of polysaccharides purified from *Hericium erinaceus* against ethanol-induced gastric mucosal injury in rats. *Int J Biol Macromol* 2020; **159**:948-56. doi: 10.1016/j.ijbiomac.2020.05.163.
 9. Liu X, Li J, Li Y, Zhang H, Zhang H. The micro mechanism and nursing of the treatment of gastric ulcer combined with gastric hemorrhage with aluminum magnesium carbonate. *Acta Microscopica* 2020; **29(3)**:1598-606.
 10. Zahid R, Akram M, Riaz M, Munir N, Shehzad M. Phytotherapeutic modalities for the management of *Helicobacter pylori* associated peptic ulcer. *Eur J Inflamm* 2020; **18**:1-16. doi.org/10.1177/2058739220968308.
 11. Jones MP. The role of psychosocial factors in peptic ulcer disease: Beyond *helicobacter pylori* and NSAIDs. *J Psychoso Res* 2006; **60(4)**:407-12. doi: 10.1016/j.jpsychores.2005.08.009.
 12. Abdulateef MM, Abdulameer RA, Muhsun LH. Prevalence of *helicobacter pylori* infection in patients with upper gastrointestinal bleeding. *Eur J Preven Med* 2019; **7(6)**: 123-9.
 13. Laporte J R, Ibanez L, Vidal X, Vendrell L, Leone R. Upper gastrointestinal bleeding associated with the use of NSAIDs. *Drug-Safety* 2004; **27(6)**:411-20. doi: 10.2165/00002018-200427060-00005.
 14. Sugisaki N, Iwakiri R, Tsuruoka N, Sakata Y, Shimoda R, Fujimoto S, et al. A case-control study of the risk of upper gastrointestinal mucosal injuries in patients prescribed concurrent NSAIDs and antithrombotic drugs based on data from the Japanese national claims database of 13 million accumulated patients. *J Gastroenterol* 2018; **53(12)**: 1253-60. doi: 10.1007/s00535-018-1483-x.
 15. Hung KW, Knotts RM, Faye AS, Pont AR, Lebwohl B, Abrams JA, et al. Factors associated with adherence to *helicobacter pylori* testing during hospitalisation for bleeding peptic ulcer disease. *Clin Gastroenterol H* 2020; **18(5)**:1091-8. doi: 10.1016/j.cgh.2019.07.037.
 16. Chen M, Wu M, Lee W, Wang H. A multiple logistic regression analysis of risk factors in different subtypes of gastric ulcer. *Hepato-gastroenterol* 2002; **49(44)**:589-92.
 17. Vestergård A, Bredahl K, de Muckadell OB, Pedersen OB, Hansen JM. Bleeding peptic ulcer. Prevalence of *Helicobacter pylori* and use of nonsteroidal anti-inflammatory drugs/ acetylsalicylic acid. *Ugeskrift for Laeger* 2009; **171(4)**:235-9.
 18. Wang JP. Clinical significance of changes of serum GAS, TNF- α and TGF- α levels after treatment in patients with peptic ulcer. *J Huaihai Med* 2009; **27(6)**:502-3.
 19. Tsai YH, Chang MH. Serum gastrin levels in children with peptic ulcer disease. *J Formos Med Assoc* 1992; **91(3)**: 283-6.
 20. Fagundes FL, Piffer GM, Périco LL, Rodrigues VP, Hiruma-Lima CA, dos Santos RC. Chrysin modulates genes related to inflammation, tissue remodeling, and cell proliferation in the gastric ulcer healing. *Int J Mol Sci* 2020; **21(3)**:760-72. doi: 10.3390/ijms21030760.
 21. Tang X, Han P, Chen Z. Changes of serum GAS, MTL, EGF AND CGRP level in patients with duodenal ulcer before and after treatment. *J Guangxi Med Univ* 1998; **15(2)**:8

• • • • •