LETTER TO THE EDITOR

OPEN ACCESS

Frequency of Keloid and Hypertrophic Scar in Post-cardiac Surgery Wounds

Sir

There are 200 cardiac operations done in our setup on monthly basis. Sternal wound complications make up about 1% of our post-cardiac surgery complications, which include superficial and deep surgical wound infection (mediastinitis and wound dehiscence). Superficial (1.1 – 6.7%) and deep wound infections (0.1 - 3.7%) are treated with dressing, debridement/re-wiring, and antibiotics.¹ Scar formation begins after the remodelling phase of wound healing. An aberrant scar formation is the most common sternal wound healing problem observed in our post-operative cardiac surgery clinics. The abnormally grown sternal scars are the only reason some patients keep visiting the post-operative clinics to get rid of them. It becomes long-term morbidity for our patients. Since the literature in cardiac surgery lacks adequate data on its prevalence so research was conducted to find out its prevalence in our post-median sternotomy wound patients.

We collected data from 300 patients to find out the frequency of keloid and hypertrophic scars in postoperative cardiac surgery patients. All coronary artery bypass grafting, valvular and aortic root replacements done *via* median sternotomy were included. There were 180 males and 120 females with ages ranging from 28 years to 66 years. People from 3 different ethnicities were taken: Sindhi, Urdu-speaking, and Pathans. All these patients had a full sternotomy extending from the jugular notch to the xiphoid process.

The frequency of hypertrophic scar came out to be 65%, keloid 10%, and normal mature scar as 25% in our post-operative cardiac surgery patients. The keloid was reported more in females as compared to males. No association of hypertrophic scar with gender was found. There was no significant difference among the three ethnic groups in the type of scar.

Scars have psychological effects. Many women have been found who developed anxiety and depression due to their scars. Astudy conducted on keloids reported that 53.3% of individuals had pain, 93% pruritis at the site of keloids, impaired quality of life in a large number of patients, and a psychological impact of 65.8%.²

There are many treatment options for hypertrophic scars and keloids. The GREMCIQ (Group for Multicenter Studies in Keloids and Hypertrophic Scars) has regarded Silicone gel sheeting as one of the widely used methods for the prevention of hypertrophic scars and keloids. It is recommended to be used soon after the suture removal when the scar is in the immature phase.^{3,4}

Prevention has always been better than cure. The patients who received topical silicone gel sheeting within 48 hours after surgery had a significantly lower occurrence of an abnormal scar than the control group (39% versus 71%).^{5,6} Currently, in cardiac surgery setups in Pakistan, none of these preventive strategies are applied as the scar is given the least importance. The purpose of this research was to bring in notice the high prevalence of aberrant scars in post-cardiac patients and to apply the widely-known strategies to reduce this morbidity in our patients.

COMPETING INTEREST:

The authors declared no competing interest.

AUTHORS' CONTRIBUTION:

WS: Substantial contribution to the concept, design of work, and collection of data.

FI: Analysis and interpretation of data.

FS: Drafting of manuscript.

All the authors have approved the final version of the manuscript to be published.

REFERENCES

- Levy AS, Ascherman JA. Sternal wound reconstruction made simple. *Plast Reconstr Surg Glob Open* 2019; 7(11):e2488. doi: 10.1097/GOX.00000000002488.
- Kassi K, Kouame K, Kouassi A, Allou A, Kouassi I, Kourouma S, et al. Quality of life in black African patients with keloid scars. *Dermatol Reports* 2020; **12(2)**:8312. doi: 10. 4081/dr.2020.8312.
- Oliveira G, Metsavaht L, Kadunc BV, Jedwab SKK, Bressan MS, Stolf HO, et al. Treatment of keloids and hypertrophic scars. Position statement of the Brazilian expert group GREMCIQ. J Eur Acad Dermatol Venereol 2021; 35(11): 2128-42. doi.org/10.1111/jdv.17484.
- Nischwitz SP, Rauch K, Luze H, Hofmann E, Draschl A, Kotzbeck P, et al. Evidence-based therapy in hypertrophic scars: An update of a systematic review. Wound Rep Reg 2020; 28(5):656-65. doi.org/10.1111/wrr.12839.
- Gold MH, Foster TD, Adair MA, Burlison K, Lewis T. Prevention of hypertrophic scars and keloids by the prophylactic use of topical silicone gel sheets following a surgical procedure in an office setting. *Dermatol Surg* 2001; **27(7)**:641-4. doi:10.1046/j.1524-4725.2001. 00356.x.
- Sidgwick GP, McGeorge D, Bayat A. A comprehensive evidence-based review on the role of topicals and dressings in the management of skin scarring. *Arch Dermatol Res* 2015; **307(6)**:461-77. doi: 10.1007/s00403-015-1572-0.

Wardah Saleem, Fayaz Iqbal and Fatima Saleem Department of Cardiac Surgery, National Institute of Cardiovascular Diseases, Karachi, Pakistan

.....

Correspondence to: Dr. Wardah Saleem, Department of Cardiac Surgery, National Institute of Cardiovascular Diseases, Karachi, Pakistan E-mail: wardah.saleem@ymail.com

Received: July 24, 2022; Revised: September 22, 2022; Accepted: October 03, 2022 DOI: https://doi.org/10.29271/jcpsp.2023.02.239

• • • • • • • • • • •