

## Origin of Kehr Sign in Traumatic Splenic Trauma

Sir,

We read with interest the case by Hashmi *et al.* of delayed splenic rupture caused by minor trauma and aim to clarify misconceptions regarding Kehr sign.<sup>1</sup> The authors used the term “Kehr’s sign” presumably in reference to hyperesthesia or pain referred to the posterior or superior left shoulder caused by blood accumulation under the left hemidiaphragm in splenic rupture or contusion. Klimpel, in a review of the literature, found a lack of compelling evidence that this sign should be attributed to Johannes Otto Kehr (1862–1916).<sup>2</sup>

In Kehr’s discussion on splenic rupture in *Handbuch der praktischen Chirurgie*, he stated that:

“There is no pathognomonic symptom for injuries to the spleen. In contusion, pain and enlargement of the spleen are typical symptoms. When rupture occurs, all the other signs and symptoms of internal bleeding are less important. In subcutaneous injury, the affected person usually experiences pain, initially on the left side, quickly spreading to involve the entire abdomen. The signs of collapse (pronounced pallor, weak pulse, feeling cold and fainting) appear. In rare cases, the injured person does not experience anything from splenic rupture, and collapse may not be present. Left shoulder pain is not common. Other manifestations involve the lungs, heart, and larynx (hoarse voice up to complete lack of voice). The latter indicates an alteration of the vagus, involving the splenic plexus through the semilunar ganglion”.<sup>3</sup>

Levy referenced Kehr’s publication on splenic injuries and left shoulder pain symptoms and presented the case of a 29-year old man with internal haemorrhage secondary to splenic injury and severe left shoulder pain, emphasising its diagnostic significance.<sup>4</sup>

Saegesser noted that phrenic nerve tenderness is found regularly in splenic rupture and subcapsular haematoma.<sup>5</sup> He described the method for eliciting this symptom as “The examination is performed by applying pressure with the thumb of the right hand from behind on the left side of the neck between the sternocleidomastoid and anterior scalene muscles and in the direction of the larynx and spine. As a control, the phrenic nerve pressure pain on the right side is checked with the same hand movement. If it is clearly detectable on the left, and together with the local findings, the assumption of a splenic injury can be assumed”.<sup>5</sup>

Confirmed is the origin of Kehr sign and the appropriate attribution to it. Both Kehr and Saegesser signs are bedside tools that may assist in diagnosing splenic rupture.

### COMPETING INTEREST:

The authors declared no competing interest.

### AUTHORS’ CONTRIBUTION:

SY, HT, EY: All involved in drafting this letter and approved the final version of the manuscript to be published.

### REFERENCES

1. Hashmi JZ, Hiraj MURK, Saleem F, Malik U, Mazari IK. Delayed splenic rupture due to an occult primary injury. *J Coll Physicians Surg Pak* 2022; **32(7)**:931-3. doi: 10.29271/jcpsp.2022.07.931.
2. Klimpel V. Stammt das Kehrsche Zeichen von Hans Kehr? Eine quellenkritische Anmerkung [Does Kehr’s sign derive from Hans Kehr? A critical commentary on its documentation]. *Chirurg* 2004; **75(1)**:80-3. doi: 10.1007/s00104-003-0796-2.
3. Kehr H. Die Chirurgie der Milz [Surgery of the spleen]. In: E. von Bergmann, P. von Burns, eds. *Handbuch der praktischen Chirurgie, Band III*. Stuttgart: Verlag von Ferdinand Enke; 1907. pp. 685-701.
4. Levy L. Zur diagnose der milzverletzungen [The diagnosis of splenic injury]. *Zentralblatt f Chir* 1910; **37(50)**:1577-1.
5. Saegesser M. Der linksseitige Phrenikusdruckpunkt ALS diagnostisches Merkmal bei Milzverletzungen [The left-sided phrenic nerve pressure point as a diagnostic feature of spleen injuries]. *Zentralblatt f Chir* 1938; **65(39)**: 2179-80.

Steven Howard Yale<sup>1</sup>, Halil Tekiner<sup>2</sup> and Eileen Scott Yale<sup>3</sup>

<sup>1</sup>Department of Internal Medicine, University of Central Florida College of Medicine, Lake Nona Blvd, Orlando, USA

<sup>2</sup>Department of the History of Medicine and Ethics, Erciyes University School of Medicine, Melikgazi, Kayseri, Turkey

<sup>3</sup>Division of General Internal Medicine, University of Florida, Florida, USA

Correspondence to: Dr. Steven Howard Yale, Department of Internal Medicine, University of Central Florida College of Medicine, Lake Nona Blvd, Orlando, USA  
E-mail: steven.yale.md@gmail.com

Received: July 19, 2022; Revised: November 07, 2022;

Accepted: November 08, 2022

DOI: <https://doi.org/10.29271/jcpsp.2023.03.360>

### AUTHOR’S REPLY

The misconceptions about the Kehr’s sign are rightly explained in the letter to the editor. Both Kehr’s and Saegesser’s signs are clinical methods specifically attributed to splenic injury. The clinicians must bear in mind that the absence of these signs

should not make them doubt their clinical judgment and complete reliance on clinical signs can result in delayed diagnosis, which can be disastrous for the patient.

Patients of blunt abdominal trauma and splenic injury should be managed according to the set guidelines. Only the gold standard investigations can put light on the nature and severity of injuries. Kehr's sign can be elicited in some patients but it has little practical application.

---

*Junaid Zia Hashmi*

.....  
*Department of General Surgery, Unit IV, Nishtar Hospital,  
Multan, Pakistan*

.....  
*Correspondence to: Dr. Junaid Zia Hashmi, Department of  
General Surgery, Unit IV, Nishtar Hospital, Multan,  
Pakistan*

*E-mail: juneehashmi@yahoo.com*

.....

