

Poppy Intoxication in Infants and Children: Hazards of a Folk Remedy

Sumera Akram, Mohammad Fazil and Kiramat Ullah

Department of Paediatrics, Mardan Medical Complex, Mardan, Pakistan

ABSTRACT

Objective: To study the clinical and laboratory profile in infants and children presented in the tertiary care hospital with poppy intake and to compare the profile between those who survived with those who died.

Study design: Observational study.

Study Place and Duration: Department Of Paediatrics, Mardan Medical Complex, Mardan, KPK, Pakistan from January 2019 to January 2020.

Methodology: All the infants and children who reported during research period with signs and symptoms (one or more) of opium poppy intoxication, i.e. meiosis, respiratory depression and decreased consciousness level along with confirmed history of giving poppy at home.

Results: A total of 32 cases of opium poppy intoxication were admitted. Their age ranged from one month to 23 months, mean age was 7.22 ± 5.43 months. Out of them, 15 (46.87%) infants and children survived, and 17 (53.13%) died, ($p < 0.001$). The children who survived had significantly shorter period of time between poppy over-dose and admission in hospital as compared to those who had died (6.0 ± 2.56 hours versus 12.47 ± 4.14 hours, $p < 0.001$). There was significantly high mortality in children who were given poppy powder mixed in water, out of 17 cases 13 died and 4 survived. Low respiratory rate, decreased oxygen saturation, aspiration pneumonia, apneic spells, cyanosis and leukocytosis >15000 cc had significant effect on mortality.

Conclusion: Poppy opium is very dangerous and hazardous for infants as it depresses respiration, causes coma; and can be fatal. People need to be educated to curb this harmful practice and authorities need to take necessary actions to stop the sale of poppy at shops and stores.

Key Words: Poppy, poisoning/Intoxication, Infants, Meiosis, Apneic spell.

How to cite this article: Akram S, Fazil M, Ullah K. Poppy Intoxication in Infants and Children: Hazards of a Folk Remedy. *J Coll Physicians Surg Pak* 2021; **31(05)**:576-581.

INTRODUCTION

The opium poppy plant *Papaver Somniferous L.* was described by Hippocrates as hypnotic, cathartic and styptic (stops bleeding of wounds) more than 350 years BC.¹ Poppies are traditional source of analgesics, ant food, anti-malarials, aphrodisiacs, rattles for baby, anti-tussives, diaphoretics, anti-diarrheal, hemostatics, hypnotics, salad vegetables, tranquilisers etc.¹ Poppy plant contains many alkaloids including morphine, codeine, narcotine, papaverine, thebaine, narcotoline and narceine.² Majority of alkaloids are in poppy milk-opium. It is prepared by drying and processing the juice extracted from fine cuts along premature poppy heads.³ It has been used for pain relief and cough. Poppy seeds are traditionally used in Pakistan as food. They normally do not contain alkaloids, but pest damage or rough harvesting can contaminate seeds with opium alkaloids.⁴

It is known that pharmacokinetics of morphine are same in adults and children.⁴ But, neonates and infants show significantly different pharmacokinetics and display longer half-life of morphine, high volume of its distribution, decreased plasma clearance and less protein binding.⁵ In addition, they also have immature blood brain barrier resulting in elevated morphine concentrations in their brain and eventually higher risk of notorious effects. After one year of age, pharmacokinetics of morphine becomes similar for both adults and neonates and infants.⁶ Because of these reasons, chances of morphine toxicity and adverse effects are higher in neonates and infants.

Generally, because of risk of serious side effects, paediatric use of morphine for chronic pain management more than three months is considered life-threatening.⁷ Morphine can also cause apnoea and seizures in neonates and infants.⁸ Other common alkaloid found in poppy is codeine. Intoxication of codeine has almost same symptoms as those of morphine, showing metabolism of codeine to morphine.⁸ Most common symptoms include respiratory depression, sleepiness, stupor, pulmonary oedema, meiosis, vomiting, cyanosis etc.⁹ In this country, people often misuse poppy for coughing and crying infants and babies, which can lead to toxicity and even death.

Correspondence to: Dr. Sumera Akram, Department of Paediatrics, Mardan Medical Complex, Mardan, Pakistan
E-mail: sumera_ak@yahoo.com

Received: February 22, 2020; Revised: October 19, 2020;

Accepted: November 27, 2020

DOI: <https://doi.org/10.29271/jcpsp.2021.05.576>

Table I: Clinical and laboratory profile of cases of poppy poisoning (n =32).

Profile			Result
Age (in months)			Mean 7.22±5.43 month Min 01 month to 23 month
Outcome	Survived Died		15 cases (46.87%) 17 cases (53.13%)
Gender	Males Females		14 (43.8%) 18 (56.3%)
Main reason of giving poppy	Fever and cough		4cases (12.5%)
	Cough		7 cases (21.9%)
	Excessive cry		9 cases 28.1%)
	Excessive cry and cough		12 cases (37.5%)
Form of giving poppy	Powder form mixed in water		15 cases (46.9%)
	Poppy pods boiled in water		17 cases (53.1%)
Time duration (hours) from intake of poppy to reach the hospital			Mean 9.44±4.75hrs Min 2 hours to max 19 hours
Clinical findings	Temperature °C		37.50±0.60° C Min 36.5°C, Max 39.1°C
	Oxygen saturation		Mean=87.3±5.82% Min=76%, Max=95%
	Meiosis		32 cases (100%)
	Low respiratory rate		27 cases (84.4%)
	Seizures		19 cases (59.4%)
	Altered sensorium		Cases 32 (100%)
	Apneic spells		18 cases (56.3%)
	Aspiration pneumonia		17cases (53.1%)
Laboratory findings	Hemoglobin ≤9gm%		20 cases (62.5%)
	Leucocytosis ≥15000/ml		21 cases (65.6%)
	Hypernatremia ≥150mEq/L		4 cases 12.5 (%)
	Hyponatremia ≤135mEq/L		1 cases (3.1%)
	Hyperkalemia >5 mEq/L		5 cases (15.6%)
	Hypokalemia <3.5 mEq/L		2cases 6.3(%)
	Serum urea >40mg%		4 cases (12.5%)
	Serum CPK ≥500 u/L		6 cases (18.8%)
Education status	Mother	Uneducated Educated	23 cases (71.9%) 9 cases (28.1%)
	Father	Uneducated Educated	20 cases (62.5%) 12 cases (37.5%)
Socioeconomic status		Poor class Middle class	24 cases (75%) 8 cases (25%)
Father job status		Jobless Employed	5 cases (15.6%) 27 cases (84.4%)
Mother job status		Housewife Employed	28 cases (87.5%) 4 cases (12.5%)
Number of siblings in the family		Four or more Less than four	17 cases (53.1%) 15 cases (46.9%)

The research has been done to study the clinical and laboratory profile in infants and children presented in the tertiary care hospitals with poppy intake and to compare the profile between those who survived with those who died.

METHODOLOGY

The study was carried out from January 2019 to January 2020 at Mardan Medical Complex, KPK, Pakistan after approval from local Ethical Committee. Informed consent was also taken from parents. The children having one or more clinical features of opium toxicity, *i.e.* meiosis, low respiratory rate, decreased level of consciousness ± seizures, vomiting along with history of intake of opium poppy at home were included in the study. The children who were administered opioids (morphine *etc.*); and neonates, whose mothers were on opioid analgesics, were excluded.

The data including age, gender, clinical presentation, reason of giving opium poppy to cases, method of opium toxicity (method of preparing and giving opium poppy to infants/children), parent's education standards, parent's employment status, socioeconomic status, number of siblings per family, time to reach hospital after intake and outcome of cases, were assessed and analysed. Education status was categorised into uneducated and educated. Socioeconomic status was categorised into poor/lower class, middle class and high class/wealthy. Employment status was categorised into unemployed and employed. For mothers, employment status was categorised into housewife and employed. All other cases of encephalitis, septic shock, seizures, drug overdose, *etc.* were excluded after careful history, examination and assessment.

SPSS version 21 was used to analyse data. Percentages was used to express frequencies, Chi-square test was used to analyse qualitative variables.

Table II: Effect of various factors on outcome in cases of poppy poisoning.

Factor	Total (N=32)	Group A (Survived) N=15	Group B (Died) N=17	p-value
Age (in months)	Mean \pm SD: 7.22 \pm 5.43	7.77 \pm 4.94	6.74 \pm 5.92	P=0.600
Gender	Male: 14 (43.75%) Females: 18 (56.25%)	5(33.33%) 10(66.67%)	9(52.94%) 8 (57.06%)	P=0.308
Main reason of giving poppy	Cough: 07 (21.87%)	6(40.0%)	1(5.88%)	P=0.005
	Excessive cry: 09 (28.12%)	6(40.0%)	3(17.64%)	
	Cough & excessive cry: 12 (37.50%)	1(6.67%)	11(64.71%)	
	Fever and cough: 4 (12.50%)	2(13.33%)	2(11.76%)	
Form of giving poppy	Poppy pods boiled in water (17 53.13%)	13(86.67%)	4(23.53%)	p<0.001
	Poppy powder mixed In water (15 46.87%)	2(13.33%)	13(76.47%)	
Time duration (hours) from intake of poppy to reach the hospital	9.44 \pm 4.75 hours	6.00 \pm 2.56 hours	12.47 \pm 4.14 hours	p<0.001
Clinical findings	Temperature °C: 37.50 \pm 0.60	37.49 \pm 0.59°C	37.51 \pm 0.63°C	p<0.001
	Oxygen saturation: 87.3% \pm 5.82	92.47 \pm 1.81%	82.82 \pm 4.07%	
	Meiosis 32 (100%)	15(100%)	17(100%)	
	Respiratory depression: 7(84.38%) Yes 5(15.62%) No	10(66.67%) 5(33.33%)	17(100%) 0(0%)	P=0.015
	Seizures: 19(59.38%) Present 13(40.62%) Absent	8(53.33%) 7(46.67%)	11(64.71%) 6(35.29%)	P=0.031
	Altered sensorium: 32(100%)	15(100%)	17(100%)	
	Apneic spells: 18(56.25%) Present 14(43.75%) Absent	5(33.33%) 10(66.67%)	13(76.47%) 4(23.53%)	
	Aspiration pneumonia: 17(53.13%) Present 15(46.87%) Absent	3(20.0%) 12(80.0%)	14(82.35%) 3(17.65%)	P=0.001
	Cyanosis: 17(53.13%) Yes 15(46.87%) No	2(13.33%) 13(86.67%)	15(88.24%) 2(11.76%)	p<0.001
	Hemoglobin \leq 9 gm% 20 (62.50%) yes 12(37.50%) No	9(60%) 6(40%)	11(64.71%) 6(35.29%)	P=0.008
Laboratory findings	Leucocytosis \geq 15000/ml 21(65.62%) Yes 11((34.38%) No	6(40%) 9(60%)	15(88.24%) 2(11.76%)	
	Hypernatremia \geq 150mEq/L 4(12.50%) Yes 28(87.50%) No	2(13.33%) 13(86.67%)	2(11.76%) 15(88.24%)	
	Hyponatremia \leq 135mEq/L 1(3.13%) Yes 31((96.87%) No	0(0%) 15(100%)	1(5.88%) 16(94.12%)	
	Hyperkalemia $>$ 5 mEq/L 5(15.63%) Yes 27((84.37%) No	1(6.67%) 14(93.33%)	4(23.53%) 13(76.47%)	
	Hypokalemia $<$ 3.5 mEq/L 2 (6.25%) Yes 30(93.75%) No	2(13.33%) 13(86.67%)	0(0%) 17(100%)	
	Serum urea $>$ 40mg% 4 (12.50%) Yes 28(87.50%) No	3(20%) 12(80%)	1(5.88%) 16(94.12%)	
	Serum CPK \geq 500 u/L 6 (18.75%) Yes 26(81.25%) No	3(20%) 12(80%)	3(17.65%) 14(82.35%)	

Table III: Other factors affecting the outcome of poppy poisoning cases.

Factor		Group A (Survived) N=15	Group B (Died) N=17	P-value
--------	--	-------------------------------	---------------------------	---------

Education mother	Uneducated 23 (71.88%)	11(73.33%)	12(70.59%)	P=1.00
	Educated 9(28.12%)	4(26.67%)	5(29.41%)	
Education father	Uneducated 20 (62.50%)	10(66.67%)	10(58.82%)	P=0.726
	Educated 12 (37.50%)	5 (33.33%)	7(41.18%)	
Mother job	Housewife 28 (87.50%)	13(86.67%)	15(88.24%)	P=1.00
	Employed 4(12.50%)	2(13.33%)	2(11.76%)	
Father job	Jobless 5 (15.63%)	2(13.33%)	3(17.65%)	P=1.00
	Employed 27 (84.37%)	13(86.67%)	14(82.35%)	
Socioeconomic status	Poor 24 (75%)	13(86.67%)	11(64.71%)	P=0.229
	Middle class 8 (25%)	2(13.33%)	6(35.29%)	
Siblings	Number of siblings ≥4 17 (53.13%)	1(6.67%)	16(94.12%)	P<0.001
	Number of siblings <4 15 (46.88%)	14(93.33%)	1(5.88%)	

Quantitative variables were expressed as mean along with standard deviation and independent t-test was used for quantitative variables. A p-value less than 0.05 was considered significant.

Normality of all quantitative variables were checked by Kolmogorov-Smirnov test; and Shapiro-wilk test was applied and variables found to be normally distributed.

RESULTS

A total of 32 cases of poppy opium toxicity came to study during the period. Their age ranged from one month to 23 months, mean age was 7.22 ± 5.43 month. There were 14 males and 18 females. Out of them, 15 (46.87%) infants and children survived, 17 (53.13%) died. All the cases were given poppy at home as a folk remedy for various purposes as shown in Table I. Most common cause of giving poppy was excessive cough and cry (12 cases), followed by excessive cry (9 cases), cough (7 cases) and fever with cough (4 cases). People gave poppy to infants and children in powder form after grinding the poppy pods and then mixing with small quantity of water (15 cases) or after boiling poppy pods in water and then feeding the infants and children (17 cases). Time duration from intake of poppy to reach hospital varied from minimum 02 hours to maximum 19 hours, mean time was 9.44 ± 4.75 hours. Mean temperature of cases was 37.50 ± 0.60 °C. Mean oxygen saturation was $87.3 \pm 5.82\%$. All the cases had meiosis (constriction of pupils) and altered sensorium. Seizures were observed in 19 cases, apneic spells in 18 cases and aspiration was seen in 17 cases, as shown in Table I.

Laboratory findings of anemia, raised leucocyte count, electrolyte imbalances, raised urea and CPK (creatinine

phosphokinase) have also been mentioned in Table II. Twenty-four cases belonged to poor socioeconomic background, and 8 belonged to middle class; however, none was from high class. Education status of parents job status of parents, and family size are shown in Table III.

All the infants, who were in critical condition, had altered sensorium and meiosis. Majority had other clinical scenarios as mentioned above. Naloxone is antidote for opium. Naloxone was given in all cases to reverse the effects of morphine alkaloid in opium. Dose of naloxone was 0.1 mg/kg. In few cases, naloxone was even repeated to reverse the effects of opium in poppy and maintain reversal. Effects of various factors on outcome (died or survived) have been assessed in Table II and III. Mean age of infants and children in group A (who survived) was 7.77 ± 4.94 months as compared to 6.74 ± 5.92 months in group B (who died), but the difference was not significant ($p=0.600$). There were 5 males and 10 females in Group A (survived) and 9 males and 8 females in Group B (who died), $p=0.308$. There was strong association between reason of poppy giving and outcome, i.e death or survival. Out of 12 cases with history of cough and excessive cry, 11 died and 01 survived; p value 0.005. However, rest of complaints, i.e cough (6 survived and 01 died), cry (6 survived and 3 died) and fever with cough (2 survived and 2 died) had more survivals as shown in Table II. There was also significantly high mortality in children who were given poppy powder mixed in water (prepared by grinding poppy pods then mixed with water), out of 17 cases 13 died and 4 survived. However, the children who were given boiled water of poppy pods had better survival (13 survived *versus* 4 died), $p<0.001$. Another important factor affecting survival was time duration to reach hospital. The mean time of children (who survived) to reach hospital was 6.0 ± 2.56 hours as compared to Group B (who died) who

had mean time of 12.47 ± 4.14 hours, $p < 0.001$. Among the clinical features, low respiratory rate, oxygen saturation, aspiration pneumonia, apneic spells and cyanosis had significant effect on mortality as shown in Table II. Among laboratory investigations, children who had leucocytosis >15000 cc had significant mortality ($p = 0.005$); however, rest of Lab findings including low hemoglobin, hyponatremia, hypernatremia, hypokalemia, hyperkalemia, raised serum urea, and raised CPK (creatinine phosphokinase) had insignificant association with outcome. Altered sensorium and meiosis were constant; therefore, p value could not be calculated for these factors.

Other factors including education status of parents, job status of parents, and socioeconomic status had no significant association with outcome, i.e. mortality versus survival. However, the infants and children who had 4 or more number of siblings, had significantly high number of deaths, i.e. 16 died and 01 survived ($p < 0.001$) as shown in Table III.

DISCUSSION

Because of easy availability of opium poppy in our country, its toxicity is not rare. Parents give poppy to infants and children in different forms for common colds, cough, crying, coryza, restlessness, abdominal colics; and it can result in overdose and even toxicity/poisoning. Opium (opiod) toxicity in children is quite serious; and without timely intervention, can cause death.^{10,11} Easy availability, cheap price and old traditional history of opium poppy are deep-rooted socially acceptable factors of using opium for minor complaints, e.g. crying etc. The presence of decreased level of consciousness, meiosis and decreased respiratory effort or apnoea should raise suspicion of opium toxicity.¹² In this study, mortality of poppy poisoning was 53.13% among infants and children. Most common cause of poppy ingestion was cough and excessive cry (seen in 37.50% of cases) and poppy was given to cases after boiling poppy pods in water (53.13) or mixing the powdered poppy in water (46.87%).

Ghaemi *et al.* reported 31 cases of opium poisoning in Iran during 2014/2015; these cases had been given opium overdose for curing cough, diarrhoea, colic and restlessness.¹¹ Zamani *et al.* had reported total 228 infants and 82 toddlers, were admitted in Lohman Hakim Hospital, Iran from 2001 to 2009 with opium toxicity. Out of these, 67 cases were result of accidental opium ingestion; however, 169 cases were overdosed consequent to folk remedy for diarrhoea, cold, cough, restlessness and fever.¹³ Accidental opium ingestion occurs when children ingest opioid drugs out of curiosity (lying within their reach due to carelessness of parents) or when parents/caregivers mistakenly give opioid instead of other drug. Imran *et al.* published local data with 22 infants of opium poisoning admitted in Lady Reading Hospital, Peshawar in 1979.¹⁴ All the cases had typical history of opium

(poppy) usage as remedy at home followed by drowsiness, cyanosis, seizures, restlessness, etc. Ira Shah reported a case of opium toxicity in India in 2012.¹⁶ The frequency of infants with opium intoxication has reduced gradually over period of time.¹⁴⁻¹⁶

In this study, educational status of parents did not have any significant effect on outcome of poppy poisoning. However, large families (>4 children) had significantly higher chances of mortality than those with less children. Water source contamination with poppy can also be a risk factor, especially in the community where water source is near poppy plants/field. People need to be educated to curb use of poppy for remedial or recreational purpose, etc. It is a serious public health issue. Authorities should play effective role by banning its cultivation and sale at stores.

CONCLUSION

Poppy-opium is given by parents to infants and children as a remedy to cure common cold, cough, crying and restlessness. Poppy opium is very dangerous and hazardous for infants and children as it depresses respiration, causes coma; and can be fatal. People need to be educated to curb this harmful practice; and authorities need to take necessary actions to stop sale of poppy at shops and stores. Proper hygiene and family planning plays an important role in decreasing morbidity and mortality of paediatric population.

ETHICAL APPROVAL:

Ethical approval was obtained from the Ethical Review Board of the Hospital prior to the start of subject research.

PATIENTS' CONSENT:

Patients' consents were obtained from the parents regarding publishing the data concerning their cases.

CONFLICT OF INTEREST:

The authors declared no conflict of interest.

AUTHORS' CONTRIBUTION:

SA: Conceived the idea, actively collected data, managed these poppy poisoning cases, and wrote the script.

MF: Collected data and managed these cases.

KU: Assisted in management of the cases and helped in data analysis & revision.

REFERENCES

1. Duke JA. Utilization of Papaver. *Economic Botany*, Vol 27, No 4, 1973. Pp 390-400. JSTOR. www.jstor.org/stable/4253448. [Accessed on Jan 20, 2020].
2. Kwiecin-Obara E, Szaponar J, Krajewska A, Witkowska A, Radoniewicz A, Szponar M. Morphine (obtained from poppy seeds) and dextrometorfan poisoning- a case report. *Przegląd Lekarski* 2015; **73(8)**: 596-8.
3. European Safety Food Authority. Opium alkaloid in poppy

- seeds: assessment updated. 16 may 2018. Accessed online 23 January 2020. <http://www.efsa.europa.eu/en/press/news/180516>
4. Martindale 2017. The complete drug reference. Morphine: Online. Pharmaceutical press, London. Available online: <http://www.medicinescomplete.com/mc/martindale/current/>.
 5. Kesawan K. Neurodevelopmental implications of neonatal pain and morphine exposure. *Pediatr Annals* 2015; 44: e260-264. doi: 10.3928/00904481-20151112-08.
 6. Lyon C, Njoku DB. Anesthetic pharmacology: physiologic states, pathophysiologic states and adverse effects. *Essentials of Pediatric Anesthesiology* 2015; 27-37. doi: 10.1017/CBO9781107375338.005.
 7. Berde CB, Walco GA, Krane EJ, Anand KJS, Arnada JV, Craig KD et al. Pediatric analgesic clinical trials designs, measures, and extrapolation: Report of an FDA scientific workshop. *Pediatrics* 2012; **129**:354-64. doi: 10.1542/peds.2010-3591.
 8. Knutson HK, Alexander J, Barregard L, Bignami M, Bruschweiler B, Ceccatelli S, et al. Update of the scientific opinion on opium alkaloids in poppy seeds. *EFSA J* 2018; **16**(5): 5243. doi: 10.2903/j.efsa.2018.5243.
 9. Dillon R, Johnston PC, Daly G. Codeine induced pulmonary edema (an unusual case of dyspnea). *QMJ* 2013; **106**: 189-290. doi: 10.1093/qjmed/hcs022.
 10. Sachdeva DK, Stadnyk JM. Are one or two dangerous? Opioid exposure in toddlers. *J Emerg Med* 2005; **29**:77-84. doi: 10.1016/j.jemermed.2004.12.015.
 11. Ghaemi N, Alikhani S, Bagheri S, Sezavar M. A cross sectional study on opioid poisoning in children at a tertiary care center. *Asia Pac J Med Toxicol* 2016; **5**:115-8. doi: 10.22038/apjmt.2016.8136.
 12. Boyer EW. Management of opioid analgesic overdose. *N Engl J Med* 2012; 367: 146-55. doi: 10.1056/NEJMr1202561.
 13. Zamani N, Sanaei-Zadeh H, Mostafazadeh B. Hallmark of opium poisoning in infants and toddlers. *Trop Doct* 2010; **40**(4):220-2. doi: 10.1258/td.2010.100134.
 14. Imran M, Uppal TB. Opium administration to infants in Peshawar region of Pakistan. *Bull Narc* 1979; **31**: 69-75.
 15. Shah I. Opium intoxication in an infant on Indian folk medicine. *J Nat Sci Biol Med* 2012; **3**(2):201-2.
 16. Trinh-Thi-Minh-HA, Nguyen-Hoc-Huong-Thu. Opium poisoning in Vietnamese infants. *South East Asia J Trop Med Public Health* 1973; **4**: 593-5.

• • • • •