Effects of Point-of-care Ultrasonography on the Diagnostic Process of Patients Admitted to the Emergency Department with Chest Pain: A Randomised Controlled Trial

Necip Gokhan Guner¹, Yusuf Yurumez¹, Murat Yucel¹, Mehmet Alacam¹, Seda Turkmen Guner² and Bilge Ercan¹

¹Department of Emergency Medicine, Sakarya University Training and Research Hospital, Turkey

²Department of Family Medicine, Sakarya University Training and Research Hospital, Turkey

ABSTRACT

Objective: To evaluate the effect of point of care ultrasonography (POCUS) performed for heart, lung, aorta, hepatobiliary and deep veins on the diagnosis, length of stay (LOS) in emergency department (ED) and cost in patients admitted to the ED with chest pain.

Study Design: Prospective randomised controlled, parallel-group trial.

Place and Duration of Study: Sakarya University Training and Research Hospital, Sakarya Turkey, from September 2018 to March 2019.

Methodology: Patients (≥ 18 years) with chest pain were randomly assigned at a 1:1 ratio to a standard diagnostic strategy (control group) or to standard diagnostic strategy supplemented with POCUS (POCUS group). Data obtained from the study were analysed using IBM SPSS Statistics 21.

Results: Two hundred and eight patients were randomly assigned to the control ($n=104$) and POCUS groups ($n=104$), respectively. The mean age was 50.42 ± 16.15 , and 54% were men. The most common comorbidity was hypertension (43%). Non-ST elevation myocardial infarction and musculoskeletal pain were the most common presumptive diagnoses. POCUS significantly reduced the LOS in ED. Detection of pathology in the POCUS increased the rate of hospitalisation. In addition, POCUS significantly shortened the LOS in the ED in patients who were discharged. The median LOS in the ED for the POCUS group was 133 min (91–279), which was significantly shorter than that of the control group at 215 min (118–372) ($p=0.006$). Although the average costs were also reduced, the difference was not statistically significant ($p=0.269$).

Conclusion: POCUS is a repeatable, practical imaging method which does not require radiation, reduces LOS in the ED statistically significant. However, further studies are needed to determine its usefulness in the ED.

Key Words: Chest pain, Cost, Emergency medicine, Length of stay, Point of care ultrasound.

How to cite this article: Guner NG, Yurumez Y, Yucel M, Alacam M, Guner ST, Ercan B. Effects of Point-of-care Ultrasonography on the Diagnostic Process of Patients Admitted to the Emergency Department with Chest Pain: A Randomised Controlled Trial. *J Coll Physicians Surg Pak* 2020; **30(12)**:1262-1268.

INTRODUCTION

Chest pain (CP) is one of the most common reasons for visiting the emergency department (ED). It is also the first symptom of many diseases.^{1,2} In the United States of America (USA), approximately 6 million patients with CP visited the ED annually, which costs approximately \$8 billion. This amount constitutes approximately 9% of all visits to the ED.³

coronary syndrome, aortic dissection, pneumothorax, pulmonary embolism, pericardial tamponade, and esophageal rupture.⁴ Important difficulties encountered in the evaluation of patients with CP in the ED include the presence of comorbidities that may cause this symptom and the need to make a differential diagnosis rapidly, but this is time-consuming and not always possible.⁵ Therefore, history and physical examination are focused on the exclusion or confirmation of serious pathology in patients visiting the ED. However, these are often insufficient, and electrocardiogram (ECG), chest X-ray, and cardiac biomarkers are often used.⁶

Another diagnostic method used in the differential diagnosis of CP is point-of-care ultrasonography (POCUS), which has become an integral method of emergency care in the USA in the last 20 years. In addition, POCUS stands out worldwide with its advanced diagnostic and therapeutic capabilities.⁷ Results of several studies in ED have shown that sonographic examina-

Correspondence to: Dr. Necip Gokhan Guner, Department of Emergency Medicine, Sakarya University Training and Research Hospital, Turkey

E-mail: gunernecipgokhan@gmail.com

Received: September 15, 2020; Revised: November 05, 2020;

Accepted: November 05, 2020

DOI: <https://doi.org/10.29271/jcpsp.2020.12.1262>

CP is often caused by life threatening diseases, such as acute

tions have excellent diagnostic accuracy for diseases most commonly encountered in patients.⁸ These diseases most often involve the heart, lungs, or deep veins of the legs, all of which can be directly visualized with sonography.⁹ The idea of using POCUS as a stethoscope is rapidly becoming integrated into clinical practice and medical education.¹⁰ Despite being a rapidly evolving diagnostic test, knowledge is needed about whether POCUS actually increases the proportion of patients who are correctly diagnosed and treated in the ED.⁸ Besides improving the health care of individuals, improved diagnostic assessment with POCUS might also have a positive effect at a societal level (by reducing health-care costs and morbidity and mortality rates) because respiratory and cardiac diseases are all major public health problems.¹¹

It was hypothesised that performing POCUS to patients presenting to the ED with CP will positively contribute to the diagnostic process, shorten the length of stay (LOS) in the ED, and reduce the average costs. The objective of this study was to evaluate the effect of POCUS performed for heart, lung, aorta, hepatobiliary and deep veins on the diagnosis, LOS in ED and costs in patients admitted to the ED with chest pain.

METHODOLOGY

This prospective, parallel-group, randomised controlled trial, with a superiority design, was conducted at the medical ED at Sakarya University Training and Research Hospital, Sakarya, Turkey. Informed consent was obtained from all individual participants included in the study. Also all procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards. The study was approved by the Sakarya University Clinical Research Ethics Committee (Decision No. 05 of 12.09.2018). The study is registered with ClinicalTrials.gov, No. NCT04429698.

Patients admitted to the ED were included in this study, if they had CP. Exclusion criteria were permanent mental disability, age <18 years, chest trauma in the last 24 hours, pregnancy, need for urgent angiography with ST elevation myocardial infarction, referral from an external center to the ED, and incomplete patient information.

The patients were randomly assigned to the POCUS or control groups. The principal investigator randomised the allocation of patients from to the groups at a 1:1 ratio by using a random number generator before the study began. When the patients admitted, it was determined in advance which numbered form was included in which group. The POCUS operator was present at alternating shifts in the ED, this situation caused inclusion of patients at different times of the day or night, and different days of the week.

Primary clinical evaluation was performed in all patients admitted to the ED. Primary clinical evaluation is defined as the evaluation of the patient by the primary physician at the time of

admission to the ED.

After the primary clinical evaluation, demographic data of all patients, characteristics of CP, location, extent, duration, vital parameters, and preliminary diagnosis of the primary physician were recorded in the study form. There was no suggestion or referral to the primary physician regarding the diagnostic approach and treatment processes.

The physician was not limited to request for complete blood count, cardiac markers, and biochemical analyses, ECG, and other imaging tests (chest X-ray, computed tomography, ultrasonography, and echocardiography), which are among the standard tests in the ED, and results can be obtained in approximately 2 hours.

At this stage, the patients were divided into two groups:

POCUS group underwent POCUS after primary clinical evaluation, in the first hour after the patients' primary clinical evaluations to evaluate the predetermined parameters in the study form for the heart, lungs, hepatobiliary, aorta, and deep veins. In the Control group; all processes and results were followed without any intervention for patients in this group, and the results were recorded in the study form.

All examinations were performed according to the American College of Emergency Physician imaging criteria by the same POCUS operator, who didn't provide primary patient care, qualified in POCUS (>250 examinations performed with focus assessed transthoracic echocardiography, >250 with focused lung ultrasonography, >250 hepatobiliary ultrasonography, >250 aortic ultrasonography and >500 with limited compression ultrasonography). Xario 100 (Toshiba, Japan) ultrasonography system was used.

After POCUS, the findings were presented to the physician in the ED who performed primary assessment, and the preliminary diagnoses, diagnostic tests, and treatment were reassessed. Without any intervention to the processes after this stage, the results, including all the processes and costs (the final amount), were recorded in the study form.

After these processes, the summaries of hospitalised patients and ED patient examination forms, laboratory, and imaging tests of the discharged patients were evaluated and recorded by a supervisor emergency medical specialist who masked POCUS findings and the patient's primary physician.

The primary endpoint was to determine the effect of POCUS on the ED LOS. The secondary endpoints were to determine the effect of POCUS on the average cost of patients, the rate of change in the preliminary diagnosis of the physician, hospitalisation and discharge rates.

For both groups, categorical endpoints were summarised as the numbers and percentages of patients with corresponding 95% confidence intervals (CIs); continuous endpoints were presented as the number of patients, mean, standard deviation, median, and interquartile range.

Table I: Demographic and clinical findings of patients in POCUS and control groups.

	POCUS group (n=104)	Control group (n=104)	All patients (n=208)
Age (years; mean $\pm$ SD)	50.28 $\pm$ 16.13	50.56 $\pm$ 16.26	50.42 $\pm$ 16.15
Gender			
Male	55 (53%)	57 (55%)	112 (54%)
Female	49 (47%)	47 (45%)	96 (46%)
Medical history			
Diabetes mellitus	25 (24%)	23 (22%)	48 (23%)
Coronary artery disease	19 (18%)	16 (15%)	35 (17%)
Hyperlipidemia	6 (6%)	7 (7%)	13 (6%)
Heart failure	4 (4%)	5 (5%)	9 (4%)
Arterial hypertension	44 (42%)	46 (44%)	90 (43%)
Stroke	1 (1%)	3 (3%)	4 (2%)
Chronic kidney disease	3 (3%)	5 (5%)	8 (4%)
Psychiatric disorders	8 (8%)	6 (6%)	14 (7%)
Chronic obstructive pulmonary disease	4 (4%)	6 (6%)	10 (5%)
Asthma	3 (3%)	3 (3%)	6 (3%)
Others	19 (18%)	12 (12%)	31 (15%)
Vital signs at admission			
Respiratory rate (breaths per min)	12 (11-12)	12 (11-12)	12 (11-12)
Saturation (%)	98 (97-99)	98 (96-98)	98 (97-99)
Systolic blood pressure (mm Hg)	120 (110-140)	120 (110-140)	120 (110-140)
Diastolic blood pressure (mm Hg)	70 (70-80)	75 (70-80)	75 (70-80)
Heart rate (beats per min)	85 (77-88)	86 (77-94)	85 (77-92)
Temperature ($^{\circ}$ C)	36.7 (36.5-36.9)	36.7 (36.5-36.9)	36.7 (36.5-36.9)
Patient reported symptoms			
Dyspnea	25 (24%)	29 (28%)	54 (26%)
Tachypnea	9 (9%)	11 (11%)	20 (10%)
Sweating	7 (7%)	15 (14%)	22 (11%)
Orthopnea	6 (6%)	5 (5%)	11 (5%)
Palpitation	10 (10%)	13 (13%)	23 (11%)
Paroxysmal nocturnal dyspnoea	4 (4%)	4 (4%)	8 (4%)
Fever	4 (4%)	2 (2%)	6 (3%)
Increase in sputum quantity or purulence	4 (4%)	2 (2%)	6 (3%)
Oedema of both legs	7 (7%)	9 (9%)	16 (8%)
Nausea	7 (7%)	12 (12%)	19 (9%)
Dyspepsia	47 (45%)	55 (53%)	102 (49%)
The fear of death	32 (31%)	22 (21%)	54 (26%)
Cough	28 (27%)	22 (21%)	50 (24%)
Dysuria	3 (3%)	1 (1%)	4 (2%)
Duration of chest pain			
Acute	87 (84%)	90 (87%)	177 (85%)
Non-acute	17 (16%)	14 (13%)	31 (15%)
Location of chest pain			
Retrosternal	16 (15%)	18 (17%)	34 (16%)
Left side of chest	54 (52%)	58 (56%)	112 (54%)
Whole chest	32 (31%)	26 (25%)	58 (28%)
Epigastric	3 (3%)	3 (3%)	6 (3%)
Back	4 (4%)	4 (4%)	8 (4%)
Radiation of chest pain			
Right shoulder and arm	- (-)	1 (1%)	1 (0%)
Left shoulder and arm	11 (11%)	10 (10%)	21 (10%)
Neck	- (-)	2 (2%)	2 (1%)
Epigastric	21 (20%)	24 (23%)	44 (21%)
Back	24 (23%)	30 (29%)	54 (26%)
Abdomen	1 (1%)	1 (1%)	2 (1%)
No radiation	51 (49%)	45 (43%)	96 (46%)
Character of chest pain			
Crushing, tightness, squeezing, pressure	25 (24%)	24 (23%)	49 (24%)
Pleuritic	59 (57%)	57 (55%)	116 (56%)
Ripping, tearing	1 (1%)	1 (1%)	2 (1%)
Sudden, sharp	17 (16%)	13 (13%)	30 (14%)
After forceful vomiting	1 (1%)	- (-)	1 (0%)
Positional	37 (36%)	45 (43%)	82 (39%)
Increasing with suppression	12 (12%)	9 (9%)	21 (10%)
Increasing with effort	20 (19%)	20 (19%)	40 (19%)
Unrelated to effort	64 (62%)	72 (69%)	136 (65%)
Burning sensation in the chest	4 (4%)	6 (6%)	10 (5%)
Imaging methods			
Electrocardiogram	100 (96%)	102 (98%)	202 (97%)
X-ray	83 (80%)	92 (88%)	175 (84%)
Computed Tomography	13 (13%)	22 (21%)	35 (17%)
Hospitalisation of patient			
Discharged	70 (67%)	74 (71%)	143 (69%)
Hospitalize	34 (33%)	30 (29%)	64 (31%)
POCUS group's ultrasound findings		Present	Absent

Cardiac ultrasound Parasternal long axis Pericardial effusion Dissection flap Pericardium thickening Valve Vegetation Aortic root >5 cm Parasternal short axis Myocardial regional wall motion abnormality Apical four chamber Right ventricle enlargement McConnell's sign Myocardial regional wall motion abnormality Systolic right atrium collapse Diastolic right ventricular collapse	4 (4%) 0 (0%) 0 (0%) 0 (0%) 0 (0%) 0 (0%) 20 (19%) 0 (0%) 0 (0%) 14 (13%) 0 (0%) 0 (0%)	100 (96%) 104 (100%) 104 (100%) 104 (100%) 104 (100%) 104 (100%) 84 (81%) 104 (100%) 104 (100%) 90 (87%) 104 (100%) 104 (100%)
Lung ultrasound B lines Pleural effusion Pneumothorax	12 (12%) 7 (7%) 4 (4%)	92 (88%) 97 (93%) 100 (96%)
Hepatobiliary ultrasound Increase in gallbladder wall thickness Stone/mud in the gallbladder Sonographic murphy's sign Enlargement of common bile duct	0 (0%) 8 (8%) 0 (0%) 0 (0%)	104 (100%) 96 (92%) 104 (100%) 104 (100%)
Abdominal aortic ultrasound Dissection flap Suprarenal aortic diameter >3 cm Infrarenal aortic diameter >3 cm	0 (0%) 0 (0%) 0 (0%)	104 (100%) 104 (100%) 104 (100%)
Ultrasound for deep vein thrombosis Thrombus in right lower extremity Thrombus in Left Lower Extremity	0 (0%) 0 (0%)	104 (100%) 104 (100%)
Data are number (%) or a mean, unless otherwise indicated. *The median values (25p-75p) were indicated because the vital signs at admission did not fit the normal distribution.		

Table II: Effects of POCUS on the differential diagnoses.

Presumptive differential diagnosis (ddx)	Pre-Pocus differential diagnosis (n=104)	Post-Pocus differential diagnosis (n=104)	p-value*
Non-ST elevation myocardial infarction	102	85	0.000
Pneumothorax	28	5	0.000
Pneumonia	32	10	0.000
Musculoskeletal pain	88	83	0.103
Pleurisy	7	6	1.000
Aort dissection	7	3	0.289
Pericarditis	16	3	0.002
Pulmonary embolism	9	0	0.004
Gastroesophageal reflux disease	16	4	0.000
Cholecystitis	4	1	0.375
Total number of DDX per patient (median (25p-75p))	3 (2-4)	2 (1,25-2)	0.000
Wilcoxon test was used for comparison of continuous endpoints, and the McNemar test as used for comparison of related groups endpoints expressed as categorically.			

The rule of three was used to calculate 95% CI in categories with no events. The Student *t* test, Mann-Whitney U-test or Wilcoxon test were used for comparison of continuous endpoints, and the χ^2 and McNemar tests were used for comparison of endpoints expressed as categorically. All tests were performed with a two-sided significance of 5%. For each endpoint, the absolute and relative effects and their corresponding 95% CIs were calculated as recommended by Altman *et al.*¹² All analyses were performed using IBM SPSS Statistics 21.

RESULTS

The demographic and clinical findings of the 208 patients included in the study are shown in Table I.

In the POCUS group, 69 pathological findings in 32 patients were detected, such as pericardial effusion, myocardial regional wall motion abnormality, gallbladder stone/sludge, B

lines, pleural effusions, pneumothorax, and aortic aneurysm. The most common of these was a myocardial regional wall motion abnormality, whereas the least common was pericardial effusion and pneumothorax (Table I).

On an average, the primary physician initially considered four differential diagnoses based on history and physical examination, and an average of two of these diagnoses were conducive to sonographic evaluation. After ultrasonography, the mean number of differential diagnoses decreased from three to two ($p<0.001$). After POCUS evaluation, a statistically significant decrease was found in most pre-diagnoses as non-ST elevation myocardial infarction (NSTEMI), pneumothorax, pneumonia, pericarditis, pulmonary embolism and gastroesophageal reflux disease ($p<0.001$; $p<0.001$; $p<0.001$; $p=0.002$; $p=0.004$; $p<0.001$, respectively, Table II).

NSTEMI and musculoskeletal pain were the most common presumed diagnoses in patients admitted to the ED.

Performing POCUS on patients presenting with non-acute CP was found to significantly decrease the LOS in the ED ($p=0.006$); in terms of pain characteristics, we found that patients in the POCUS group who described pleuritic pain significantly wait less than the control group ($p=0.003$).

The use of ECG and chest X-ray and imaging methods together with POCUS in patients presenting to the ED with CP has reduced the LOS in the ED.

When the last diagnoses were evaluated, POCUS significantly reduced the LOS in the ED in patients with the final diagnosis of musculoskeletal pain. After the patients were examined and subsequently discharged, the LOS in the ED in the POCUS group was shorter than that in the control group. When all patients were considered, we found that performing POCUS significantly shortened the LOS in the ED. In addition, performing POCUS did not reduce the average costs (Table III).

When the POCUS group was examined, the LOS in the ED was prolonged in patients with pathology detected during ultrasonography, and the average costs were higher (respectively, 212 min (113-391) vs. 123 min (85-257) $p=0.012$; 30.9 ± 18.4 vs. 22.6 ± 13.1 $p=0.026$). In addition, the presence of pathology on ultrasonography significantly increased hospitalization rates (discharged = 7 patients 21.9% vs. hospitalized = 25 patients 78.1%, $p<0.001$).

DISCUSSION

CP is among the most common causes of referral to the ED and is among the most important symptoms of cardiovascular diseases, which ranks first among the most common causes of deaths worldwide. CP is caused by many cardiac and non-cardiac causes, and that serious increases in mortality and morbidity occur when the approach is not performed correctly and in time, increase the importance of diagnostic approaches. In this context, the effectiveness of POCUS was investigated to ensure that patients with CP visiting the ED were diagnosed and treated early, as well as to shorten the LOS in the ED and reduce the diagnostic process.

Studies by Lamsam *et al.* and Buhumaid *et al.* investigated the diagnostic efficacy of POCUS in patients presenting to the ED with CP and respiratory symptoms.^{13,14} However, to the best of authors' knowledge, no randomized controlled clinical trial has investigated the effect of POCUS on the LOS in ED and costs in patients with CP.

The demographic characteristics of the patients in this study were consistent with those of previous studies in the literature related to CP. For example, in the study by Lamsam *et al.*, the mean age was 59 years, and in our

study, it was 50.42 years.¹⁴ In the study by Buhumaid *et al.*, 55% of the patients with CP who visited the ED were men. Similarly, this rate was 57.6% and 64% in the study by Lamsam *et al.* and Carubbi *et al.*, respectively. In our study, similar to the literature, 54% of men with CP visited the ED more frequently than women with CP.¹³⁻¹⁵

POCUS, a diagnostic tool recommended in the differential diagnosis approach, is very useful in making quick and accurate decisions, revealing pathologies and shaping pre-diagnoses. Using POCUS, pericardial effusion, myocardial contraction defect, enlarged heart chambers, B lines in the lung, pleural effusion, pneumothorax, aortic aneurysm, aortic dissection, deep vein thrombosis, pathologies of the gallbladder, and many organ pathologies can be detected.⁷ Considering the studies conducted to investigate pathologies, the detection rate was 20% and 30% in the study by Yates *et al.* and Buhumaid *et al.*, respectively.^{13,16} The most frequent pathological finding is decreased ejection fraction of the heart due to myocardial wall motion defect, and the most common irregular pleural line and B lines in terms of lung. The detection rate of pathological findings in our study was 34%. The most common pathological findings were myocardial wall motion defect and diffuse B lines in the lung. These results are also consistent with the literature.^{13,16}

It is stated in ACEP's ultrasonography guide that POCUS can have a positive effect on the diagnosis and treatment process in patients with CP admitted to the ED.⁷ In this study, after performing ultrasonography, the median number of differential diagnoses decreased from three (2-4) to two (1.25-2.00; $p<0.001$). Following POCUS evaluation, a statistically significant decrease was found in all sonographically identifiable diagnoses. These results are consistent with previous studies in which POCUS was found to be a feasible and accurate diagnostic tool in patients presenting with CP.^{13,17} In addition, the detection of pathology in the POCUS has increased the rate of hospitalization similar to the study by Elikashvili *et al.*¹⁸

In our study, the median value of the LOS in the ED of the POCUS and control groups was 133 (91-279) and 215 (118-372) minutes, respectively, which was statistically significant ($p=0.006$). A study investigating the effect of POCUS on the LOS in the ED in patients with CP admitted to the ED has not been found in the literature, but the studies by Lin *et al.* investigating patients with soft tissue infection, by Park *et al.* for patients with renal colic, by Elikashvili *et al.* for patients with suspected appendicitis, and by Wilson *et al.* achieved the same result in patients admitted to the ED due to pelvic pain and vaginal bleeding.¹⁸⁻²¹

Table III. Effects of POCUS on the LOS and average costs.

	Pocus group *los (min) median (25p-75p)	Control group *los (min) median (25p-75p)	p-value**	Pocus group average costs (mean $\pm$ sd ; \$)	Control group average costs (mean $\pm$ sd ; \$)	p-value**
--	---	---	-----------	--	--	-----------

Duration of chest pain						
Acute	160 (98-290)	184 (108-375)	0.544	22.1 ± 10.7	26.7 ± 13.5	0.307
Non-acute	130 (89-270)	222 (119-375)	0.006	25.7 ± 16.0	27.8 ± 17.9	0.418
Character of chest pain						
Pressure, tightness, Pleuritic	328 (174-448)	368 (218-450)	0.779	28.8 ± 14.6	27.2 ± 11.0	0.662
	114 (82-219)	195 (101-321)	0.003	24.5 ± 16.2	30.2 ± 21.1	0.106
Positional	120 (92-232)	185 (115-280)	0.040	26.1 ± 18.6	26.3 ± 17.9	0.975
Unrelated to efor	123 (89-278)	185 (102-324)	0.075	23.2 ± 13.3	25.8 ± 15.8	0.304
Pre-diagnosis						
NSTEMI	133 (92-281)	215 (117-363)	0.011	25.2 ± 15.4	26.5 ± 15.8	0.559
Musculoskeletal pain	126 (88-265)	196 (111-309)	0.016	23.5 ± 14.4	25.7 ± 16.1	0.358
Pneumothorax	123 (86-274)	225 (120-311)	0.083	27.1 ± 19.9	38.7 ± 30.0	0.161
Pneumonia	222 (121-345)	242 (133-403)	0.459	29.6 ± 18.7	34.4 ± 15.5	0.307
Imagining methods						
ECG	133 (91-274)	215 (117-375)	0.006	24.6 ± 14.8	26.7 ± 15.8	0.337
X-ray	130 (92-269)	225 (120-372)	0.002	26.0 ± 16.3	27.8 ± 17.5	0.485
CT	155 (103-444)	357 (215-490)	0.082	51.3 ± 22.6	46.4 ± 21.3	0.529
Diagnosis						
NSTEMI	328 (214-438)	362 (223-410)	0.855	28.1 ± 14.8	29.9 ± 14.3	0.682
Musculoskeletal pain	110 (81-222)	182 (96-275)	0.015	20.4 ± 10.7	21.5 ± 11.0	0.576
Hospitalisation of patient						
Discharged	110 (82-211)	185 (102-313)	<0.001	21.4 ± 12.3	24.3 ± 13.2	0.171
Hospitalise	279 (135-413)	280 (134-443)	0.721	32.9 ± 17.9	36.0 ± 22.9	0.557
All patients	133 (91-279)	215 (118-372)	0.006	25.2 ± 15.3	27.7 ± 17.3	0.269

*LOS: Length of stay; **p-values calculated using Mann Whitney-U and independent sample T-test. Data are number median, unless otherwise indicated.

In addition, in this study, POCUS did not affect the rate of discharge or hospitalisation, whereas the discharged patients in POCUS group significantly had shorter LOS in the ED [POCUS group: 110 (82-211); control group: 185 (102-313) minutes, $p < 0.001$]. The authors believe that this result will decrease the density of the ED and improve the quality of care in clinics, such as our clinic, with a high number of ED visits.

The limitations of this study are: it is a single-center study design used for this trial, and one physician performed all the sonographic examinations; therefore, the results might not be applicable to other EDs or physicians performing POCUS. Patients could not be consecutively enrolled when the investigator for POCUS was not present in the ED, and this might have introduced bias. Involving several physicians performing sonographic examinations would have been preferable, but this was not possible because there were no other emergency physicians with ultrasonography qualification when the study began. Another limitation of this study was that no attempt to ensure the accuracy of the ultrasound-aided decision-making with regard to the final diagnosis, and no blinded evaluation of the images to ensure accuracy of the image interpretation.

CONCLUSION

POCUS is a reproducible, irradiated, practical imaging method performed by ED physicians and successfully detects pathological findings and helps in the diagnostic process by narrowing the differential diagnosis range and by reducing the LOS in the ED. Crowded ED may improve the delivered quality of patient care. However, cost reduction was not seen. The results obtained should be supported with additional studies because there is no previous study on CP.

ETHICAL APPROVAL:

The study protocol was approved by the local Ethics Committee [IRB No:16214662/050.01.04/74].

PATIENTS' CONSENT:

Informed consents were obtained from all participants.

CONFLICT OF INTEREST:

Authors declared no conflict of interest.

AUTHORS' CONTRIBUTION:

NGG: Led and conceived the project, and authored the manuscript and statistics.

YY: Data collection, compiling and discussion.

MY: Contributed to design articles, collected and analysed data.

MA: Contributed to collect and analyse data.

STG: Collected and analysed data, literature review and discussion.

BE: Helped perform the analyses with constructive discussions.

REFERENCES

- Hamm CW, Goldmann BU, Heeschen C, Kreyman G, Berger J, Meinertz T. Emergency room triage of patients with acute chest pain by means of rapid testing for cardiac troponin T or troponin I. *N Engl J Med* 1997; **337**(23): 1648-53. doi: 10.1056/NEJM199712043372302.
- Solinas L, Raucci R, Terrazzino S, Moscariello F, Pertoldi F, Vajto S, et al. Prevalence, clinical characteristics, resource utilization and outcome of patients with acute chest pain in the emergency department. A multicenter, prospective, observational study in North-Eastern Italy. *Ital Hear J* 2003; **4**(5):318-24.
- Diercks DB, Hollander JE. Acute coronary syndromes. In: Judith E. Tintinalli O, John Ma, Donald Yealy, Garth D. Meckler J. Stephan Stapczynski, David M. Cline SHT, Ed. Tintinalli's emergency medicine: A comprehensive study guide. 9th edition. New York: McGraw-Hill Education 2019; p.334-52.
- Mahler SA. Chest Pain. In: Judith E. Tintinalli, O. John Ma, Donald Yealy, Garth D. Meckler, J. Stephan Stapczynski, David M. Cline SHT, Ed. Tintinalli's emergency medicine a

- comprehensive study guide. 9th edition. New York: *McGraw-Hill Education* 2019; p.329-34.
5. Kontos MC, Diercks DB, Kirk JD. Emergency department and office-based evaluation of patients with chest pain. In: Mayo clinic proceedings. *Elsevier Ltd* 2010; **85(3)**:284-99. doi: 10.4065/mcp.2009.0560.
 6. Niska R, Bhuiya F, Xu J. National hospital ambulatory medical care survey: 2007 emergency department summary. *Natl Health Stat Report* 2010; **26(26)**:1-31.
 7. Ultrasound guidelines: Emergency, point-of-care and clinical ultrasound guidelines in medicine. Vol. 69, *Annals of Emergency Medicine*. Mosby Inc.; 2017. p. e27-54.
 8. Laursen CB, Sloth E, Lassen AT, Christensen R de P, Lambrechtsen J, Madsen PH, et al. Point-of-care ultrasonography in patients admitted with respiratory symptoms: A single-blind, randomised controlled trial. *Lancet Respir Med* 2014; **2(8)**:638-46. doi: 10.1016/S2213-2600(14)70135-3.
 9. Ray P, Birolleau S, Lefort Y, Becquemin MH, Beigelman C, Isnard R, et al. Acute respiratory failure in the elderly: Etiology, emergency diagnosis and prognosis. *Crit Care* 2006; **10(3)**:10(3):R82. doi: 10.1186/cc4926.
 10. Moore CL, Copel JA. Point-of-care ultrasonography. *N Eng J Med* 2011; **364(8)**:749-57. doi: 10.1056/NEJMra0909487.
 11. Kröger K, Küpper-Nybelen J, Moerchel C, Moysidis T, Kienitz C, Schubert I. Prevalence and economic burden of pulmonary embolism in Germany. *Vasc Med* 2012; **17(5)**: 303-9. doi: 10.1177/1358863X12449363.
 12. Robert G. Newcombe DGA. In: Altman D, Machin D, Bryant T GM, Ed. *Statistics with Confidence: Confidence Intervals and Statistical Guidelines*. 2nd Edition. London: Wiley; 2000. p. 45-7.
 13. Buhumaid RE, St-Cyr Bourque J, Shokoohi H, Ma IWY, Longacre M, Liteplo AS. Integrating point-of-care ultrasound in the ED evaluation of patients presenting with chest pain and shortness of breath. *Am J Emerg Med* 2019; **37(2)**:298-303. doi: 10.1016/j.ajem.2018.10.059
 14. Lamsam L, Gharahbaghian L, Lobo V. Point-of-care ultrasonography for detecting the etiology of unexplained acute respiratory and chest complaints in the emergency department: A prospective analysis. *Cureus* 2018; **10(8)**: e3218. doi: 10.7759/cureus.3218.
 15. Carubbi C, Masselli E, Pozzi G, Mattioli M, Martini S, Goldoni M, et al. Combination of Platelet expression of PKCepsilon and cardiac troponin-I for early diagnosis of chest pain patients in the emergency department. *Sci Rep* 2019; **9(1)**:2125. doi: 10.1038/s41598-019-38624-5.
 16. Yates J, Royse CF, Royse C, Royse AG, Canty DJ. Focused cardiac ultrasound is feasible in the general practice setting and alters diagnosis and management of cardiac disease. *Echo Res Pract* 2016; **3(3)**:63-9. doi: 10.1530/ERP-16-0026.
 17. Ahn JH, Jeon J, Toh HC, Noble VE, Kim JS, Kim YS, et al. SEARCH 8Es: A novel point of care ultrasound protocol for patients with chest pain, dyspnea or symptomatic hypotension in the emergency department. *PLoS One* 2017; **12(3)**:e0174581. doi: 10.1371/journal.pone.0174581.
 18. Elikashvili I, Tay ET, Tsung JW. The effect of point-of-care ultrasonography on emergency department length of stay and computed tomography utilization in children with suspected appendicitis. *Acad Emerg Med* 2014; **21(2)**: 163-70. doi: 10.1111/acem.12319.
 19. Lin MJ, Neuman M, Rempell R, Monuteaux M, Levy J. Point-of-Care Ultrasound is Associated With Decreased Length of Stay in Children Presenting to the Emergency Department With Soft Tissue Infection. *J Emerg Med* 2018; **54(1)**: 96-101. doi: 10.1016/j.jemermed.2017.09.017.
 20. Park YH, Jung RB, Lee YG, Hong CK, Ahn J-H, Shin TY, et al. Does the use of bedside ultrasonography reduce emergency department length of stay for patients with renal colic? a pilot study. *Clin Exp Emerg Med* 2016; **3(4)**:197-203. doi: 10.15441/ceem.15.109
 21. Wilson S. Point-of-care ultrasound versus radiology department pelvic ultrasound on emergency department length of stay. *World J Emerg Med* 2016; **7(3)**:178. doi: 10.5847/wjem.j.1920-8642.2016.03.003.

• • • • •