Effect of Anastomosis Technique on Pancreatic Fistula Formation in Proximal Pancreaticoduodenectomy

Ozgun Yildirim Yildirim¹ and Nazmi Özer²

¹Izmir Katip Celebi Universitesi, Izmir, Turkey

²Adana City Research and Training Hospital, Adana, Turkey

ABSTRACT

Objective: To find out the association between the pancreatic fistula development and the pancreaticoduodenectomy anastomosis technique performed in three different ways; namely, telescopic pancreatojejunostomy (PJ), ducto-mucosal Wirsung jejunostomy (WJ), and Peng's variant of the telescopic technique.

Study Design: A descriptive study.

Place and Duration of Study: Izmir Katip Celebi Universitesi Hospital, Turkey, from January 2011 to January 2018.

Methodology: A total of 144 proximal pancreaticoduodenectomy procedures were performed at the study centre. Patients' data was obtained from the outpatient files and hospital information system, retrospectively; and divided into three groups according to the pancreatic anastomosis techniques. All three groups were compared in terms of eight parameters. Preoperative parameters were age, gender, and serum direct bilirubin values; whereas, postoperative parameters were pathology, pancreatic fistula, drain amylase, serum albumin value and early mortality.

Results: All variables by anastomosis type belonging to three groups differed for fistula rates ($p=0.384$), pathology types ($p=0.142$), preoperative bilirubin ($p=0.631$) and postoperative albumin ($p=0.516$) levels, early mortality ($p=0.242$) parameters without reaching statistical significance ($p>0.05$); however, the low anastomosis leak rates in Peng's technique were remarkable. POPF (post-operative pancreatic fistula) developed in 34% patients under 60 years vs. 17.6% patients over 60 years of age ($p=0.029$); and 13.7% females vs. 29.9% of male patients ($p=0.032$).

Conclusion: No factors other than age and gender were found to be significant alone in the development of pancreatic fistula after proximal pancreaticoduodenectomy.

Key Words: Postoperative pancreatic fistula, Pancreaticoduodenectomy, Pancreatojejunostomy, Telescopic pancreatojejunostomy (PJ), Ducto-mucosal Wirsung jejunostomy (WJ), Peng's telescopic technique.

How to cite this article: Yildirim OY, Özer N. Effect of Anastomosis Technique on Pancreatic Fistula Formation in Proximal Pancreaticoduodenectomy. *J Coll Physicians Surg Pak* 2020; **30(05)**:480-484. DOI: <https://doi.org/10.29271/jcpsp.2020.05.480>.

INTRODUCTION

Proximal pancreaticoduodenectomy (PPD, Whipple) operation is a procedure performed in pancreas head cancers, duodenum, distal choledochal and ampulla tumors, chronic pancreatitis, and also major injuries of the pancreatic head containing the portal vein.¹ It is a complex and high-risk surgical procedure.² In recent years, the mortality rate has decreased markedly in the last two decades.³ Appropriate patient selection, increased surgical experience, developments in preoperative and postoperative follow-up, and multidisciplinary approach have contributed to better results. Nevertheless, even in experienced centres, mortality is still 1-5% and total morbidity is 30-50%.^{1,4,5}

The most important complication after PPD is the development of pancreatic fistula which may lead to other major complications.

According to ISGPF (International Study Group on Postoperative Pancreatic Fistula), the incidence of POPF varies between 5% and 30%.^{6,7}

Different types of pancreatojejunostomy anastomosis are used in PPD surgery. The commonly used techniques are: pancreatojejunostomy, Wirsung jejunostomy, and Peng's technique. Although these techniques have individual or double comparisons in terms of fistula, there are no studies comparing all 3 techniques in literature.⁸⁻¹⁰

The aim of this study was to determine the association between the pancreatic fistula and three different anastomosis techniques in patients undergoing proximal pancreaticoduodenectomy.

METHODOLOGY

This descriptive study was carried out by retrospectively analysing the data of 144 patients who underwent proximal pancreaticoduodenectomy (PPD) between January 2011 and January 2018 with the approval of the ethical committee number 24.03.2018/63 in General Surgery Clinic of Izmir Katip Celebi University Hospital.

Patient's age, gender and serum direct bilirubin values were recorded in the preoperative period.

Correspondence to: Nazmi Özer, Adana City Research and Training Hospital, Adana, Turkey
E-mail: nazmiozer83@gmail.com

Received: April 11, 2020; Revised: May 08, 2020;

Accepted: May 13, 2020

DOI: <https://doi.org/10.29271/jcpsp.2020.05.480>

It was examined for pancreatic fistula development status, pathology of the resected tissue, serum albumin value, early mortality and anastomosis technique, in the postoperative period. The data were obtained by examining the medical records including pre- and post-operative clinical notes, surgical reports, laboratory results and pathology reports. Patients who were operated with a technique other than the three studied anastomotic techniques were excluded.

POPF (postoperative pancreatic fistula) was defined as being that the amylase value in the drain fluid obtained on the third postoperative day or after is three times higher than the serum amylase value, as suggested by the ISGPF (international study group on postoperative pancreatic fistula).

Three reconstruction techniques were evaluated to ensure the continuity of pancreatic stump namely end-to-end invagination (telescopic) pancreatojejunostomy (PJ, Figure 1a); end-to-side ducto-mucosal (Wirsungojejunostomy, WJ, Figure 1b); and Peng's technique, a variant of the telescopic technique (Figure 1c). Stent was used in telescopic PJ anastomosis group only and none of the patients was given octreotide in postoperative period. The patients were divided into two groups as those who developed and not developed pancreatic fistula after PD operation.

Figure 1: The varieties of pancreatoenteric anastomosis (a) Telescopic PJ (b) Ductomucosal WJ (c) Its schemes Peng's technique.
(em: everted mucosa, rp: residue of pancreas)

SPSS 22.0 programme was used for analysis of data. The normality of data distribution was evaluated with the Shapiro-Wilk test. One-Way ANOVA test was used together with bootstrap results in comparison of more than two independent groups by quantitative variables. In comparison of categorical data, Pearson, Chi-Square test was tested with Monte Carlo Simulation technique and Fisher's exact test. Quantitative data were expressed as mean $\pm$ SD (standard deviation) and range (maximum-minimum) values in the tables. In addition, categorical data were expressed in n (number) and percentages (%). Values with $p < 0.05$ were considered significant.

RESULTS

The mean age of the patients was 62.94 (18-86). 38.2% of the patients were aged 60 and under, and 61.8% of them were over the age of 60. It was observed that POPF developed in 34% of patients aged 60 and under, and 17.6% of them over 60 years old. The differences between two groups were statistically significant ($p=0.029$).

Majority *i.e.* 62.5% of the patients were males and 37.5% were females. POPF developed in 13.7% female patients and 29.9% male patients. The differences between two groups were statisti-

cally significant ($p=0.032$).

Table I: The relationship between risk factors and POPF status developing in patients underwent PPD.

POPF	Absent (n=105) n (%)	Present (n=33) n (%)	p-value
Age			
≤60	35 (33.3)	18 (54.5) ^b	0.029
>60	70 (66.7)	15 (45.5)	2.40 (1.08-5.32) ^a
Gender			
Woman	44 (41.9)	7 (21.2)	0.032
Man	61 (58.1)	26 (78.8) ^b	2.68 (1.07-6.72) ^a
Pathology			
Adenocarcinoma	86 (81.9)	24 (72.7)	0.514
Intraepithelial neoplasia	5 (4.8)	3 (9.1)	
Invasive ductal carcinoma	0 (0.0)	1 (3.0)	
Chronic pancreatitis	3 (2.9)	2 (6.1)	
Non-specific duodenitis	1 (1.0)	0 (0.0)	
Neuroendocrine carcinoma	6 (5.7)	1 (3.0)	
Pseudopapillary Tm.	3 (2.9)	1 (3.0)	
Villous adenoma	1 (1.0)	1 (3.0)	
Drain amylase			
Normal	16 (100.0)	0 (0.0)	<0.001
High	0 (0.0)	33 (100.0)	
Serum direct bilirubin (preoperative)			
<0.5	16 (100.0)	13 (39.4)	0.655
≥0.5	0 (0.0)	20 (60.6)	
Serum albumin (postoperative)			
≤3	84 (89.4)	27 (96.4)	0.454
>3	10 (10.6)	1 (3.6)	
Ex in the early period			
Absent	103 (98.1)	31 (93.9)	0.242
Present	2 (1.9)	2 (6.1)	

Pearson Chi-Square Test (Monte Carlo) - Fisher Exact Test (Exact): comparison of categorical data.
*Odds ratio (95% Confidence interval) : age and gender/popf; *Reference for Odds ratio: age and gender/popf.

Overall, 34.72% of the patients were anastomosed with the telescopic PJ, 59.03% of them with the ducto-mucosal WJ, 6.25% of them with the Peng's technique. The POPF rate was determined as 19.1% (9/47) in the telescopic technique, 28% (23/82) in the ducto-mucosal technique, and 11.1% (1/9) in the Peng's technique. In 3 patients, each operated by telescopic technique and by ductomucosal technique, leakage status was evaluated as uncertain due to drain and patient problems. The pathological diagnoses of the resected specimen showed 80.6% adenocarcinoma, 5.6% intraepithelial neoplasia, 4.9% neuroendocrine carcinoma, 3.5% chronic pancreatitis, 2.8% pseudopapillary tumors, 1.4% villous adenoma, and 0.7% each of invasive ductal carcinoma and nonspecific duodenitis ($p=0.514$).

Regarding postoperative serum albumin levels, it was 3.0 g/dL; and below in 90.98% (111/122), 24.3% of these patients developed POPF. The patients with postoperative serum albumin levels above of 3.0 g/dL were 8.9% (11/122); and 9.1% of these patients developed POPF ($p=0.454$). Preoperative serum direct bilirubin value of 0.5 mg/dL and above were found in 57.2% (79/138), 25.3% of these patients developed POPF. The same level was below 0.5 mg/dL in 42.7% (59/138), and 22.0% of these patients developed POPF ($p=0.655$).

No statistically significant difference was found between the development pancreatic fistula, the highness of serum bilirubin

Table II: Relationship between anastomosis techniques and risk factors in patients undergoing PPD.

Anastomosis type	Ductomucosal WJ. (n=85) n (%)	Peng's (n=9) n (%)	Telescopic PJ. (n=50) n (%)	Total (N=144) n (%)	p-value
Age Mean $\pm$ SD. (max-min)	62.17 $\pm$ 11.23 (86-18)	61.89 $\pm$ 10.74 (76-42)	64.46 $\pm$ 11.37 (83-33)	62.94 $\pm$ 11.23 (86-18)	0.500
≤ 60 >60	34 (40.0) 51 (60.0)	3 (33.3) 6 (66.7)	18 (36.0) 32 (64.0)	55 (38.2) 89 (61.8)	0.856
Sex Woman Man	31 (36.5) 54 (63.5)	2 (22.2) 7 (77.8)	21 (42.0) 29 (58.0)	54 (37.5) 90 (62.5)	0.533
Pathology Adenocarcinoma Intraepithelial neoplasia Invasive ductal carcinoma Chronic pancreatitis Non-specific duodenitis Neuroendocrine carcinoma Pseudopapillary Tm. Villous adenoma	74 2 1 2 0 3 3 0	5 2 0 1 0 0 1 0	37 4 0 2 1 4 0 2	116 (80.6) 8 (5.6) 1 (0.7) 5 (3.5) 1 (0.7) 7 (4.9) 4 (2.8) 2 (1.4)	0.142
Fistula status Absent Present	59 (72.0) 23 (28.0)	8 (88.9) 1 (11.1)	38 (80.8) 9 (19.1)	105 (76.1) 33 (23.9)	0.384
Drain amylase Normal High	7 (23.3) 23 (76.7)	3 (75.0) 1 (25.0)	6 (40.0) 9 (60.0)	16 (32.7) 33 (67.3)	0.077
Serum direct bilirubin (preoperative) <0.5 ≥ 0.5	37 (43.5) 48 (56.5)	5 (55.6) 4 (44.4)	19 (38.0) 31 (62.0)	61 (42.4) 83 (57.6)	0.631
Serum albumin (postoperative) ≤ 3 >3	66 (89.2) 8 (10.8)	7 (100.0) 0 (0.0)	40 (93.0) 3 (7.0)	113 (91.1) 11 (8.9)	0.516
Ex in the early period Absent Present	81 (95.3) 4 (4.7)	8 (88.9) 1 (11.1)	46 (92.0) 4 (8.0)	135 (93.7) 9 (6.3)	0.535

One Way ANOVA Test (Method: Bootstrap): Used together with bootstrap results in comparison of more than two independent groups by quantitative variables. Pearson Chi-Square Test (Monte Carlo): Comparison of categorical data. SD: Standard Deviation, Max: Maximum, Min: Minimum.

level in the preoperative period and the lowness of serum albumin level in the postoperative period. While the early mortality rate of patients undergoing PPD was 6.3% (9/144), when the groups developing POPF and those not developing were examined, no statistically significant difference was found between the presence of fistula and early mortality rates ($p = 0.242$, Tables I and II).

DISCUSSION

Many risk factors are described for pancreatic fistula development after PPD.¹¹ Age, gender, degree of jaundice, malnutrition, pathology of periampullary region disease, resected pancreatic part, consistency of pancreatic stump, width of pancreatic duct, operation time, type of resection, technique of pancreatic anastomosis and intraoperative blood loss are included among these risk factors.^{7,11} In this study, the rate of POPF in women over 60 years and female gender was statistically lower than male under 60 years of age.

Herrera *et al.* defined the risk factors for POPF. That factors are: over 70 years, male gender, and soft pancreatic tissue.¹²

In the study by Kow *et al.*, PPD operation was performed in patients with over 65 years and under. When results were evaluated, the rate of POPF and complications were higher in patients over 65 years.¹³ While, the gender parameter in this study is parallel with the literature, the age parameter is not parallel with the literature. The likely reason for this result is additional disease and physical performance score, which may be better in elderly patients than the younger group.

In the present series, POPF was found in 22.9% (33/144); whereas mortality rate was 6.3% (9/144). The POPF rate was similar, and mortality rate was lower, when compared to the literature. In the series by Duffas *et al.*, mortality was 11% and the rate of POPF was 18%;¹⁴ whereas Fang *et al.* determined that postoperative mortality was 8.9%, postoperative morbidity was 56.4% and POPF was 17.6%.¹⁵ The lower mortality rate after resection in the present series may be related not to the surgeon case load but to the level of experience in the various departments involved in the detection and management of postoperative complications, such as gastroenterology, radiology, or intensive care.

While 9 (19.1%) of 47 patients, who underwent telescopic PJ, developed POPF, 23 (28%) of 82 patients developed POPF who underwent ductomucosal WJ. In the Peng's technique, this rate was found to be 11.1% (1/9). Although the comparison of these three methods is not statistically significant ($p = 0.384$), the low rate of fistula in Peng's technique was remarkable compared to other techniques.

Yang *et al.* decided to choose the anastomosis technique according to the state of the residual pancreatic tissue. They concluded that ductomucosal WJ should be performed without a stent in patients whose pancreatic duct diameter is dilated ($>3\text{mm}$). Telescopic type PJ should be performed with stent in patients whose pancreatic residual tissue is soft and pancreatic duct diameter is not dilated, again, no statistically significant difference was found when anastomosis types, telescopic PJ and ducto mucosal WJ were compared in terms of POPF.¹⁶

In the study by Yingbin *et al.*, POPF rates were found in 100 patients, upon whom they performed the Peng's technique.¹⁷ In the study by Javier *et al.*, 2 (6.7%) POPF was determined in 30 patients who underwent Peng's technique.¹⁸ In the light of the available literature and the results of this study, leakage rates are lower in Peng's technique compared to other techniques.

Two general factors-serum direct bilirubin level in the preoperative period and serum albumin level in the postoperative period-were not statistically significant, when compared between groups with and without pancreatic fistula. These results are in line with the literature.^{16,19} In their first series, Yeo and Cameron regarded these two factors among risk factors in pancreatic fistula development.¹ Whereas, in their later study, they reported them as a general risk factor, not specific to pancreatic fistula.¹⁹

In the prevention of pancreatic fistula, methods such as placing a stent in the pancreatic duct and the use of octreotide in the postoperative period have also been tried; accordingly, it was not observed that the stenting had effects such as protection from fistula and reduction in rate of leak, and also it was observed that it caused different complications such as pancreatic duct obstruction.^{20,21} In addition, the use of octreotide was not observed to be a prohibitive on fistula development with reducing pancreatic exocrine secretion.^{19,22-25}

Considering the pathology results, adenocarcinoma was observed in 80.6%. Additionally, pancreatitis and other tumoral formations were observed at a lower frequency. When the fistula rates were compared in relation with the pathology results, there was no statistically significant difference.

CONCLUSION

In this study, no statistically significant difference was found between the three techniques in terms of POPF rates. Leakage rate was lower in Peng's technique compared to other techniques. No effect of parameters except age and gender was found statistically significant on POPF rates being lower in female gender and over age 60 patients. Ducto mucosal WJ

should be performed in patients with hard pancreatic tissue and dilated duct. Peng's technique should be performed in patients with soft pancreatic tissue and non-dilated/narrowed duct.

ETHICAL APPROVAL:

The ethics committee approval of our study was obtained before the study started and it was accepted with the approval of the ethics committee at the General Surgery Clinic of Izmir Katip Celebi University Atatürk Training and Research Hospital.

PATIENTS' CONSENT:

Consent for the participation in study was not obtained from patients as data was collected from medical record without disclosing the identity of participants.

CONFLICT OF INTEREST:

Authors declared no conflict of interest.

AUTHORS' CONTRIBUTION:

NO: Concept, supervision, interpretation and critical review.

OYY: Design, data collection, processing and writing manuscript.

REFERENCES

1. Yeo CJ, Cameron JL, Sohn TA, Lillemoe KD, Pitt HA, Talamin MA, *et al.* Six hundred and fifty consecutive pancreaticoduodenectomies in the 1990s: Pathology, complications, and outcomes. *Ann Surg* 1997; **226(3)**:248-57.
2. Kayahara M, Nagakawa T, Ueno K, Ohta T, Takeda T, Miyazaki I. An evaluation of radical resection for pancreatic cancer based on the mode of recurrence as determined by autopsy and diagnostic imaging. *Cancer* 1993; **172(7)**:2118-23.
3. Gouma DJ, van Geenen RC, van Gulik TM, de Haan RJ, de Wit LT, Busch OR, *et al.* Rates of complications and death after pancreaticoduodenectomy: Risk factors and the impact of hospital volume. *Ann Surg* 2000; **232(6)**:786-95.
4. Yeo CJ. The Johns Hopkins experience with pancreaticoduodenectomy with or without extended retroperitoneal lymphadenectomy for periampullary adenocarcinoma. *J Gastrointest Surg* 2000; **4(3)**:231-2.
5. Pedrazzoli S. Pancreatoduodenectomy (PD) and postoperative pancreatic fistula (POPF): A systematic review and analysis of the POPF-related mortality rate in 60,739 patients retrieved from the English literature published between 1990 and 2015. *Medicine (Baltimore)* 2017; **96(19)**:e6858.
6. Ho IG, Kim JK, Hwang HK, Kim JY, Park JS, Yoon DS. Does international study group on pancreatic fistula (ISGPF) classification need modification after distal pancreatectomy? *Korean J Hepatobiliary Pancreat Surg* 2014; **18(3)**:90-3.
7. Distler M, Kersting S, Rückert F, Kross P, Saeger H.D, Weitz J, *et al.* Chronic pancreatitis of the pancreatic remnant is an independent risk factor for pancreatic fistula after distal pancreatectomy. *BMC surgery* 2014; **14**:54.
8. Kawaida H, Kono H, Hosomura N, Amemiya H, Itakura J, Fujii H, *et al.* Surgical techniques and postoperative management to prevent postoperative pancreatic fistula after pancreatic surgery. *World J Gastroenterol* 2019; **25(28)**:3722-37.
9. Wang W, Zhang Z, Gu C, Liu Q, Liang Z, He W, *et al.* The optimal choice for pancreatic anastomosis after pancreaticoduodenectomy: A network meta-analysis of

- randomized control trials. *Int J Surg* 2018; **57**:111-6.
10. Gupta S, Attri AK, Sharma R, Gureh M, Nasir MI. Binding pancreaticojejunostomy: Is It Safe? *Niger J Sur* 2020; **26(1)**: 48-52.
 11. Crippa S, Salvia R, Falconi M, Butturini G, Landoni L, Bassi C. Anastomotic leakage in pancreatic surgery. *HPB (Oxford)* 2007; **9(1)**:8-15.
 12. Herrera J, Zazpe C, Sánchez P, Tarifa A, Eguaras I, Lera JM. Feasibility and morbidity and mortality in two hundred consecutive cases of pancreaticogastrostomy after pancreaticoduodenectomy. *Cir Esp* 2019; **97(9)**:501-9.
 13. Kow AWC, Sadayan NA, Ernest A, Wang B, Chan C.Y, Ho CK, et al. Is pancreaticoduodenectomy justified in elderly patients? *Surgeon* 2012; **10(3)**:128-36.
 14. Hu BY, Wan T, Zhang WZ, Dong JH. Risk factors for postoperative pancreatic fistula: Analysis of 539 successive cases of pancreaticoduodenectomy. *World J Gastroenterol* 2016; **22(34)**:7797-805.
 15. Duffas JP, Suc B, Msika S, Fournat G, Muscari F, Hay JM, et al. Associations for research in surgery. A controlled randomized multicenter trial of pancreatogastrostomy or pancreato-jejuno-stomy after pancreatoduodenectomy. *Am J Surg* 2005; **189(6)**:720-9.
 16. Yang YM, Tian XD, Zhuang Y, Wang WM, Wan YL, Huang YT. Risk factors of pancreatic leakage after pancreaticoduodenectomy. *World J Gastroenterol* 2005; **11(16)**: 2456-61.
 17. Liu Y, Wang J, Xu B, Li H, Zhe T, Fang H, et al. Peng's pancreaticojejunal anastomosis (binding pancreaticojejunostomy) for the pancreato-duodenectomy- A new procedure ensuring no leakage. *Chinese-German J Clin Oncol* 2002; **1**:65-7.
 18. Targarona J, Barreda L, Pando E, Barreda C. Is peng's pancreaticojejunal anastomosis more effective than mucosa-mucosa anastomosis in duodenopancreatectomy for pancreatic and peri-ampullary tumours? *Cirugía Española (English Edition)* 2013; **91(3)**:163-8.
 19. Yeo CJ, Cameron JL, Lillemoe KD, Sauter PK, Coleman J, Sohn TA, et al. Does prophylactic octreotide decrease the rates of pancreatic fistula and other complications after pancreaticoduodenectomy? Results of a prospective randomized placebo-controlled trial. *Ann Surg* 2000; **232(3)**:419-29.
 20. Casadei R, Ricci C, Silvestri S, Campra D, Ercolani G, D'Ambra M, et al. Peng's binding pancreaticojejunostomy after pancreaticoduodenectomy. An Italian, prospective, dual-institution study. *Pancreatology* 2013; **13(3)**:305-9.
 21. Suzuki Y, Fujino Y, Tanioka Y, Hiraoka K, Takada M, Ajiki T, et al. Selection of pancreaticojejunostomy techniques according to pancreatic texture and duct size. *Arch Surg* 2002; **137(9)**: 1044-7.
 22. Poon RT, Lo SH, Fong D, Fan ST, Wong J. Prevention of pancreatic anastomotic leakage after pancreaticoduodenectomy. *Am J Surg* 2002; **183(1)**:42-52.
 23. Peng YP, Zhu XL, Yin LD, Zhu Y, Wei JS, Wu JL, et al. Risk factors of postoperative pancreatic fistula in patients after distal pancreatectomy: A systematic review and meta-analysis. *Sci Rep* 2017; **7(1)**:185.
 24. Yang X, Patel S, Cunningham SC. The harm of prophylactic octreotide in pancreatoduodenectomy: More artefact than fact? *HPB (Oxford)* 2015; **17(4)**:371.
 25. Jin K, Zhou H, Zhang J, Wang W, Sun Y, Ruan C, et al. Systematic review and meta-analysis of somatostatin analogues in the prevention of postoperative complication after pancreaticoduodenectomy. *Dig Surg* 2015; **32(3)**: 196-207.

• • • • •